



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

0 253 455
A1

12

EUROPEAN PATENT APPLICATION

21 Application number: **87201379.2**

22 Date of filing: **20.07.87**

51 Int. Cl.: **C 12 N 15/00, C 12 N 1/20,**
C 12 N 1/16, C 12 P 21/00,
A 61 K 37/02, C 07 K 13/00,
A 61 K 39/395, G 01 N 33/68,
G 01 N 33/577
// (C12N1/20,C12R1:125),
(C12N1/20,C12R1:10),
(C12N1/20,C12R1:07)

30 Priority: **18.07.86 GB 8617594**
23.12.86 GB 8630699

43 Date of publication of application: **20.01.88**
Bulletin 88/3

64 Designated Contracting States: **AT BE CH DE ES FR GB**
GR IT LI LU NL SE

71 Applicant: **GIST-BROCADES N.V., Wateringseweg 1,**
NL-2611 XT Delft (NL)

72 Inventor: **Van Ooyen, Albert Johannes Joseph, Laan van**
Holdenburg 23, NL-2271 AR Voorburg (NL)
Inventor: **Andreoli, Peter Michael, Molenlaan 111,**
NL-3055 EJ Rotterdam (NL)
Inventor: **Van Mourik, Jan Aart, Maraboestraat 1,**
NL-1171 SX Badhoevedorp (NL)
Inventor: **Pannekoek, Hans, Stommeerweg 36,**
NL-1431 EW Aalsmeer (NL)

74 Representative: **Huygens, Arthur Victor, Dr. et al, c/o**
Gist-Brocades N.V. Patent & Trademarks Department
Wateringseweg 1 PO Box 1, NL-2600 MA Delft (NL)

54 Method for the preparation of proteins with factor VIII activity by microbial host cells; expression vectors, host cells, antibodies.

57 Microbial systems are provided for producing peptides having biological and physiological factor VIII activity, in being immunologically cross-reactive with factor VIII and physiologically active in providing for clotting activity. The protein products vary from immunologically cross-reactive fragments to proteins comprising the two Factor VIII subunits joined together directly or by a bridge. High yields of the products are obtained in isolatable and biologically active form.

EP 0 253 455 A1

ACTORUM AG

BEST AVAILABLE COPY

S-2428

13 juli 1987

METHOD FOR THE PREPARATION OF PROTEINS WITH FACTOR VIII
5 ACTIVITY BY MICROBIAL HOST CELLS; EXPRESSION VECTORS, HOST
CELLS, ANTIBODIES.

INTRODUCTION

10 Technical Field

The invention relates to a microbiological process for the preparation of proteins with factor VIII activity.

Background

15 Factor VIII (also referred to as factor VIIIc) is a medically important protein since it is an essential component of the blood coagulation mechanism. It occurs in the blood plasma of normal healthy individuals at a concentration of about 100 ng/ml. Factor VIII deficiency results in a bleeding
20 disorder called hemophilia A, which occurs mainly in males because the factor VIII gene is located on the X-chromosome. The disease is the major inherited bleeding disorder, occurring in about 0.01% of the male population.

Hemophilia A can be treated by administering factor
25 VIII containing blood plasma obtained from healthy people. This treatment, however, has several disadvantages. Firstly, the supply of factor VIII is limited and very expensive because the concentration in the blood of donors is very low. Secondly, the yields of current plasma fractionation methods
30 are low. In fact, there is a shortage of factor VIII. Thirdly, every hemophilia A patient receives samples collected from a large number of donors which leads to a high risk of acquiring infectious diseases, such as those caused by hepatitis non-A, non-B, hepatitis B or AIDS viruses present in donor blood.
35 Another problem is the development of antibodies against factor VIII preparations in so-called inhibitor patients.

These problems have provoked an intense search for alternative methods for the production of factor VIII, which avoid the use of donor blood, viz. by recombinant DNA techniques and three groups have reported molecular cloning of 5 factor VIII cDNA obtained from mRNA.

International Patent Application WO 85/01961 (Genetics Institute) describes the preparation of cDNA coding for human factor VIII or for subunits with factor VIII 10 activity. The cDNA is cloned, whereafter a mammalian COS-7 cell line is transformed with a vector containing said cDNA.

European patent application EP-A-160457 (Genentech) describes a method of transfecting a mammalian (Baby Hamster Kidney) cell line with a vector containing DNA coding for 15 functional human factor VIII.

European Patent Application EP-A-150735 (Chiron) describes a method for the preparation of the factor VIII protein and of fragments thereof with factor VIII activity. The vector containing DNA coding for said protein came to 20 expression in mammalian cell lines, particularly in COS cells (Truett et al., 1985).

Although the above-mentioned publications state that microorganisms could be used as the host cells in an expression system, no experimental results describing this are 25 presented, leading to the inevitable conclusion that the reduction to practice of this procedure has encountered unexpected difficulties at the experimental level.

The cited literature shows that proteins with factor VIII activity can be prepared by recombinant DNA techniques, 30 but only in mammalian cell lines. However, the concentration of factor VIII produced by the cell lines was very low; in fact, too low for commercial production. An additional disadvantage of the use of these cell lines is that they may be derived from tumor material and thus may contain substances 35 which are hazardous to health.

It was known moreover, e.g. from European

applications EP-A-150735 (Chiron) and EP-A-123945 (Scripps) and from Brinkhous et al., (1985) that factor VIII activity should be attributed not only to the protein factor VIII itself, but also to certain proteolytic cleavage products of it. Thrombin is a proteolytic enzyme which, when present in the blood, partially digests factor VIII into subunit polypeptides two of which have molecular weights of 92 kDa and 80 kDa. It is believed that when a complex of these two subunit proteins or parts thereof (Faye et al., 1986, Eaton et al., 1986) appear in the blood the factor VIII activity is enhanced.

DEFINITIONS

Factor VIII is the protein which corrects the clotting defect known as hemophilia A. Factor VIII circulates in the blood combined with a larger protein, the von Willebrand factor (vWf), which is believed to protect the sensitive factor VIII against early degradation.

Factor VIII (human) is characterized by the amino acid sequence as described in Table I. The numbering of the sequence starts with Ala(1), the first amino acid after the signal sequence of 19 amino acids. The last amino acid is Tyr(2332). This numbering is used throughout the specification.

In the literature, factor VIII is frequently reported as the combination of factor VIIIC, which for the purposes of this invention will be referred to as factor VIII and factor VIIIVWf, the von Willebrand factor. In this invention, factor VIII is the combination of a 80 kDa subunit and a 92 kDa subunit which are derived from a much larger precursor protein, which will be referred to as the factor VIII precursor. In the precursor, the 80 kDa subunit and the 92 kDa subunit are joined by a bridge, which is highly glycosylated.

Fragments of factor VIII will normally include at

least a portion of one of the two subunits and may include, but will normally not include, portions of the precursor protein outside of the 80 kDa and 92 kDa subunits.

In referring to a peptide having factor VIII biological activity, immunological cross reactivity with one of the subunit proteins will be sufficient to fulfill this requirement.

"Pharmaceutical preparations with factor VIII activity" means any preparation which can correct the clotting defect associated with factor VIII deficiency. The correction of the clotting defect may be effected by direct action of the recombinantly prepared protein, by providing one or both of the 80 kDa and 92 kDa subunits or functionally equivalent analogs thereof, or by neutralizing antibodies against factor VIII protein present in inhibitor patients so that a subsequently administered factor VIII protein, analog or functionally equivalent fragment is able to display its full activity.

For convenience, the term "protein" will be used to include proteins and smaller polypeptides.

SUMMARY OF THE INVENTION

The present invention provides novel constructs for producing proteins having factor VIII biological and physiological activity in isolatable form. The production processes employ microbial hosts, particularly yeast or bacteria, transformed with an expression cassette comprising a transcriptional and translational initiation region which efficiently functions in the host and an open reading frame encoding the protein of interest. Preferred constructs encode a signal leader for secretion of the encoded product, as well as a fused protein which includes a bridge between the two subunits.

Expression may be achieved employing extrachromosomal elements or with integration of the expression cassette into the host genome.

BRIEF DESCRIPTION OF THE DRAWINGS

5 FIGURE 1 : ANALYSIS OF PURIFIED FACTOR VIII-VON WILLEBRAND
FACTOR COMPLEX (PANEL A) AND PURIFIED FACTOR VIII
POLYPEPTIDES DEVOID OF VON WILLEBRAND FACTOR (PANEL
B) BY IMMUNOBLOTTING.

10 Factor VIII containing samples were subjected to SDS-
polyacrylamide gel electrophoresis. After separation, proteins
were transferred to nitrocellulose sheets and, after
appropriate washing steps, incubated with antibodies to factor
VIII polypeptides.

15

A: Lane 1: polyclonal antibody RH 63271

Lane 2: polyclonal antibody RH 63272

Lane 3: polyclonal antibody RH 63275

Lane 4: monoclonal antibody CLB-CAg 9

20 Lane 5: monoclonal antibody CLB-CAg 65

B: Lane 6: monoclonal antibody CLB-CAg 21

Lane 7: monoclonal antibody CLB-CAg 62.

Panel A and B are run for different times. Arrows indicate
25 positions of 80 kDa and 92 kDa proteins.

FIGURE 2 : STRATEGY FOR SYNTHESIS OF A cDNA CLONE TO FACTOR
VIII mRNA.

30

Specific primers, complementary to the indicated
parts of factor VIII mRNA were used for priming cDNA
synthesis. A second oligonucleotide which can also be used for
priming of the next cDNA fragment, is used for detection of
35 correct cDNA clones. Nucleotide sequences of primers used are
as follows:

- 1: 3'GGAAGGTAGACGGACTGGGGAAG 5'
- 2: 3'GGACTCCTACCTCCAAGACCCG 5'
- 3: 3'CCTTTAGGGTTCTCTTCAGTGG 5'
- 4: 3'GGTAGTCTGTAAACCGTCGTCC 5'
- 5 5: 3'CGTACTCCACCGTATGACCATG 5'
- 6: 3'GCCACTCGACGGACACCTGCG 5'

Resulting cDNA fragments are subcloned with restriction enzyme sites present in circles: E: EcoRI; B: BamHI; P: PstI; Sa: 10 SacI. Numbers below fragments refer to the size of the corresponding fragment in base pairs. Plasmids containing fragments A,B,C,D and E are called pCLB81, pCLB82, pCLB83, pCLB84 and pCLB85, respectively.

15 FIGURE 3 : A MAP OF pOL5-21, AS AN EXAMPLE OF A
pOL5-n TYPE PLASMID





- Plasmids of the pOL5-n type consist of:
-  : B. licheniformis T5 DNA sequences;
 - 20  : Chymosin-encoding sequences;
 -  : Alpha-amylase promoter, ribosome binding site, ATG initiation codon and the DNA encoding 29 amino acids of the alpha-amylase signal sequences;
 - 25 * { : DNA encoding n amino acids of the mature alpha-amylase protein (in the case of pOL5-21, 21 amino acids).
 -  : Vector pUB110 with a Bacillus origin of replication (Bac.ori) and the gene for neomycin resistance (Nm).
 - 30 m.c.s. : multicloning sites.

FIGURE 4 : CONSTRUCTION OF pOL5-DELTA PLASMIDS

In plasmid pOL5-21delta the B. licheniformis DNA 35 sequences upstream the alpha-amylase gene have been deleted. Symbols as in Figure 3.





FIGURE 5A AND 5B : CONSTRUCTION OF pOL92 PLASMIDS.

5 5A : pOL92-T-n plasmids are derived partly from pOL5-n
(Figure 3) and pGB842 which is the SalI-EcoRV
fragment containing the 92 kDa subunit DNA from
pOL92-A-n plasmids are derived partly from pOL5-n
and the SalI-ApaI fragment containing the 92 kDa
10 subunit DNA from pGB842.

Symbols used are:

- B. ori: Bacillus origin of replication.
- E. ori: E. coli origin of replication
- 15 - Ap: the pAT153 gene encoding ampicilline resistance
- Nm: the pUB110 gene encoding neomycin resistance
- pA: the gene expression and signal sequences of the
→ B. licheniformis alpha-amylase gene
- m.c.s.: multicloning sites

20

-  : B. licheniformis T5 DNA sequences
-  : Factor VIII 92 kDa-encoding cDNA insert
-  : Kluyveromyces lactis lactase terminator(T)
-  : Chymosin DNA

25

5B : Alternatively, pOL92-A-n plasmids can be
constructed from pOL92-T-n plasmids and pOL-5-n as
depicted for pOL92-A-21.

30

FIGURE 5C: CONSTRUCTION OF pOL92-DELTA PLASMIDS

In plasmid pOL92-T-21delta the B. licheniformis DNA
35 sequences upstream the alpha-amylase gene have been deleted.
Symbols used as in Figure 5A,B.

FIGURE 6 : POLYACRYLAMIDE GEL ANALYSIS OF CELL LYSATES
OF B. SUBTILIS TRANSFORMANTS.

The samples were analysed on a 6% SDS-polyacrylamide gel and
5 stained with Coomassie brilliant blue.

Lane 1: cell lysate from Bacillus subtilis transformed with
pOL92-T-21 Lane 2: cell lysate from Bacillus subtilis
transformed with pOL-5-21


10 Lane 3: molecular weight marker

The arrow indicates the protein band of 88 kDa in pOL92-T-21.

FIGURE 7: STRATEGY FOR DELETION OF THE B. LICHENIFORMIS ALPHA-
15 AMYLASE STRUCTURAL GENE

Plasmid pGB33 contains a 3.3kb EcoRI fragment
harbouring the B. licheniformis alpha-amylase gene and
flanking DNA sequences.

20 In plasmid pGB33delta the whole alpha-amylase gene
has been deleted. Symbols used are:

 : alpha-amylase encoding sequences

NmS : neomycin sensitive phenotype

25 NmR : neomycin resistant phenotype

Amy⁺ : alpha-amylase proficient phenotype

Amy⁻ : alpha-amylase deficient phenotype

Other symbols as in Figure 3.

30 FIGURE 8: POLYACRYLAMIDE GEL ANALYSIS OF CELL LYSATES OF
B. LICHENIFORMIS TRANSFORMANTS.

lanes 1-4 contain extracellular proteins and lanes 6-9
intracellular proteins. Staining with Coomassie brilliant blue.

35








- lane 1: Bacillus licheniformis T5-16 transformed with pOL92-A-21
- lane 2: Strain T5-16, control
- lane 3: Strain T5-16, control
- 5 lane 4: Strain T5-16, transformed with pOL92-T-21
- lane 5: Molecular weight markers
- lane 6: Bacillus licheniformis T5-16 transformed with pOL92-A-21
- lane 7: Strain T5-16, control
- 10 lane 8: Strain T5-16, control
- lane 9: Strain T5-16, transformed with pOL92-T-21

The arrow indicates the 92 kDa subunit protein.

15 FIGURE 9: CONSTRUCTION OF pOL80 PLASMIDS

Plasmid pOL80-T-40delta has been constructed from pGB852, M13mpl8-80 kDa and pOL5-40delta as depicted.

20 Symbols used are:

-  : lactase-encoding DNA
 : lactase terminator
 : lactase promoter
 : G418 resistance marker
25  : remaining part from factor VIII B-domain
 : 3' non-coding region of factor VIII cDNA
 : 80 kDa protein encoding DNA

Other symbols as in Figure 3.

30 FIGURE 10: DETECTION OF FACTOR VIII POLYPEPTIDES PRODUCED BY TRANSFORMED BACILLUS STRAINS WITH SPECIFIC MONO- AND POLYCLONAL ANTIBODIES

- A. Immunoblot detection of the 92 kDa subunit protein with
- 35 monoclonal antibody CLB-CAg9 and goat anti-rabbit alkaline phosphatase.

Lanes 1-5 contain proteins from culture supernatants. Lane 1 and 2: B. licheniformis T9 transformed with pOL92-T-2ldelta and pOL92-T-2delta, respectively. Lane 3: B. licheniformis T9 strain, control. Lanes 4 and 5: B. licheniformis T9 transformed with pOL5-2ldelta and pOL5-2delta, respectively. Lane 6 shows the pattern obtained with a factor VIII cryoprecipitate. Lane 7: high molecular weight blot protein markers from BRL (Bethesda). Arrow at left indicates the position of recombinant 92 kDa (r92 kDa) subunit. Arrow at right indicates the position of plasma-derived 92 kDa (pd92 kDa) subunit.

B. Immunoblot detection of the 80 kDa subunit protein with polyclonal antibody RH63272 and goat anti-rabbit alkaline phosphatase.

Lanes 3-7 contain proteins from culture supernatants. Lane 1: high molecular weight blot protein markers from BRL (Bethesda). Lane 2 shows the pattern obtained with a factor VIII cryoprecipitate. Lane 3: B. licheniformis T9 transformed with pOL5-40delta vector. Lanes 4-7: B. licheniformis T9 transformed with pOL80-T-40delta, pOL80-T-2ldelta, pOL80-T-10delta and pOL80-T-2delta, respectively. Arrow at left indicates the position of plasma-derived 80 kDa (pd80 kDa) subunit doublet. Arrows at right indicate the position of recombinant 80 kDa (r80 kDa) subunit and a cleavage product of r80 kDa.

C. Immunoblot detection of the 80 kDa subunit protein with polyclonal antibody RH 63272 and goat anti-rabbit alkaline phosphatase.

Lanes 2-7 contain proteins from cell lysates. Lane 1: shows the pattern obtained with a factor VIII cryoprecipitate. Lanes 2, 4 and 6: B. subtilis 1A40 transformed with pOL80-T-40delta, pOL80-T-2ldelta and pOL80-T-2delta, respectively. Lanes 3, 5 and 7: B.

subtilis 1A40 transformed with pOL5-40delta, pOL5-21delta and pOL5-2 delta, respectively. Arrow at right indicates the position of plasma derived 80 kDa (pd80 kDa) subunit doublet. Arrow at left indicates the position of recombinant 80 kDa (r80 kDa) subunit.






D. Immunoblot detection of a 92kDa/80kDa fusion protein with polyclonal antibody RH 63272 and goat anti-rabbit alkaline phosphatase.

Lanes 1-8 contain proteins from cell lysates. Lanes 1-4: controls of B. subtilis 1A40 transformed with pPROM3-4 vector. Lanes 5-8: four independently isolated colonies of B. subtilis 1A40 transformed with pPROM92/80PB. Arrow at right indicates the position of the 92 kDa-80 kDa fusion protein.

FIGURE 11: CONSTRUCTION OF pPROM92/80PB PLASMID

Plasmid pPROM92/80PB has been constructed from pGB861 and pPROM3-4 as depicted.

Symbols used are:

-  : alpha-amylase encoding sequences
-  : 92 kDa and 80 kDa-encoding DNA
-  : remaining part from factor VIII B-domain
-  : B. licheniformis T5 DNA sequences
-  : B. licheniformis 3-4 promoter

Other symbols as in Figure 9.

FIGURE 12: STRUCTURE OF pGBdLAC.

Plasmid pGBdLAC contains an XbaI fragment harbouring the K. lactis lactase gene and promoter and terminator sequences. The

middle fragment between two *Cla*I sites has been deleted from the gene.




FIGURE 13: STRUCTURE OF pGB850.

10

Plasmid pGB850 contains the 92 kDa subunit-encoding DNA under control of the lactase promoter. Symbols used are:





FIGURE 14: STRUCTURE OF pGB851

20


Plasmid pGB851 contains the major part of the 92 kDa subunit DNA with the bovine prochymosin gene in phase at the *Bam*HI site.



Other symbols as in Fig. 13.

30 FIGURE 15: STRUCTURE OF pGB852

Plasmid pGB852 contains the 80 kDa-encoding DNA fused to the lactase gene.



encoding DNA.

===== : 3' non-coding region of factor VIII cDNA

Other symbols as in Fig. 13.

5

FIGURE 16: STRUCTURE OF pGB853

Plasmid pGB853 contains the 80 kDa-encoding DNA under control of the lactase promoter.

10

Symbols as in fig. 15.

FIGURE 17: STRUCTURE OF pGB861

15 Plasmid pGB861 contains a factor VIII cDNA from which part of the B-domain has been removed.

===== 92 kDa and 80 kDa-encoding DNA

||||| remaining part from B-domain

20

Other symbols as in fig. 15.

FIGURE 18: RNA ANALYSIS OF K. LACTIS TRANSFORMED WITH pGB850 AND pGB852

25 A: lane 1: Control K. lactis

2: K. lactis transformed with pGB850

3: Independent K. lactis transformed with pGB850

B. lane 1: Control K. lactis

30 2: K. lactis transformed with pGB852

3: Independent K. lactis transformed with pGB852

Arrows indicate the position of ribosomal RNA bands.

35 FIGURE 19: DETECTION OF 92 kDa AND 80 kDa SUBUNIT-CONTAINING PROTEINS PRODUCED BY TRANSFORMED K. LACTIS STRAINS WITH SPECIFIC MONO- AND POLYCLONAL ANTIBODIES

- 5 A. Immunoblot detection of the lactase-80 kDa fusion protein
with polyclonal antibody RH 63275 and donkey anti-rabbit
peroxidase. Lanes 1 and 3: cell extract of the parental
K. lactis strain made in the presence of 0.05% Tween 80 or
2% NP40, respectively. Lanes 2 and 4: cell extract of
K. lactis expressing the lactase-80 kDa fusion protein,
made in the presence of 0.05% Tween 80 or 2% NP40,
10 respectively. Lane 5 shows the pattern obtained with a
factor VIII preparation. Arrow indicates the position of
the fusion protein.
- 15 B. Immunoblot detection of the lactase-80 kDa fusion protein
with polyclonal antibody RH 63272 and goat anti-rabbit
alkaline phosphatase. Lanes 1 and 3: cell extract of the
parental K. lactis strain made in the presence of 0.05%
Tween 80 or 2% NP40, respectively. Lanes 2 and 4: cell
extract of K. lactis transformed with pGB852 and expressing
20 the lactase-80 kDa fusion protein, made in the presence of
0.05% Tween 80 or 2% NP40, respectively. Lane 5 shows the
pattern obtained with a factor VIII preparation. Arrow
indicates the position of the fusion protein.
- 25 C. Immunoblot detection of the lactase-80 kDa fusion protein
with monoclonal antibody CLB-CAg 65 and goat anti-mouse
alkaline phosphatase. All cell extracts were made in the
presence of 0.05% Tween 80. Lanes 1-3: three independently
isolated strains transformed with plasmid pGB850 containing
30 the coding sequence for the 92 kDa subunit; lane 4 shows a
factor VIII preparation; lanes 5-8 independently isolated
strains transformed with plasmid pGB852 containing the
coding sequence for the lactase-80 kDa fusion; lane 9:
parental K. lactis strain. Arrow indicates the position of
35 the fusion protein.

- D. Immunoblot detection of the 92 kDa-prochymosin fusion with an antiserum against chymosin and donkey anti-rabbit peroxidase. Lane 1: parental K. lactis strain; lanes 2-4: independently isolated clones transformed with plasmid pGB851 containing the 92 kDa-prochymosin coding sequence; lane 5: yeast transformed with pGB850 containing the 92 kDa coding sequence only; lane 6: yeast transformed with pGB852 containing the 80 kDa coding sequence; lane 7: transformed yeast producing prochymosin; lane 8 shows the pattern of a commercial chymosin preparation. The cell extracts investigated in lanes 9-12 were incubated at pH 2 before preparing the samples. Lane 9: parental yeast strain; lanes 10-12: independently isolated clones transformed with the 92 kDa-prochymosin coding sequence. Arrow at left indicates position of prochymosin fusion. Arrows at right are at position of prochymosin and chymosin, respectively.
- E. Immunoblot detection of the 80 kDa subunit protein with polyclonal antibody RH 63272. Lane 1: K. lactis transformed with pGB852 (cf Fig. 19B). Lanes 2-5: Independent K. lactis colonies transformed with pGB853. Lane 6: control strain.
- F. Immunoblot detection of the 92 kDa sununit protein with monoclonal antibody CLB-CAg9. Lane 1: K. lactis transformed with pGB850. Lane 2: Control K. lactis.
- G. Immunoblot detection of the 92 kDa and 80 kDa subunits expressed simultaneously in one cell. Lanes 1,2 and 3: detection of 80 kDa protein with polyclonal antibody RH 63272 Lane 1 control lane, lanes 2 and 3, two different transformed strains. Lanes 4,5 and 6 detection of 92 kDa protein with monoclonal antibody CLB-CAg9. Lane 4: control lane; lanes 5 and 6, same strains as in lanes 2 and 3.
- H. Immunoblot detection of the 92 kDa-80 kDa fusion protein

with polyclonal antibody RH 63272. Lane 1: Control K. lactis. Lanes 2-4: Independent K. lactis colonies transformed with pGB 861.

- 5 I. Immunoblot detection of secretion of the 92 kDa-80 kDa fusion protein with polyclonal antibody RH 63272. Lane 1: Plasma factor VIII. Lanes 2 and 3: cell supernatant and pellet of untransformed K. lactis. Lanes 4 and 5: cell supernatant and pellet of transformed K. lactis.

10

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

In accordance with the subject invention, novel DNA
15 constructs are provided which include expression cassettes having an open reading frame encoding at least a portion of either or both the 80 kDa or 92 kDa subunits of factor VIII. The constructs will usually be prepared stepwise employing appropriate vectors, with cloning at each of the stages and
20 analysis of the product to ensure that the proper organization and sequences have been obtained. The resulting constructs may then be introduced into appropriate vectors, particularly shuttle vectors, for transformation into preferred hosts and expression of the desired protein. The product may be
25 expressed within the cytoplasm or may be secreted into the nutrient medium or a combination of both, where the unmaturred protein which retains the signal leader may be further processed by appropriate enzymatic treatment. The resulting products are found to have biological and in many cases
30 physiological activity, in being able to be used to promote clotting activity, either by themselves or in combination with endogenous factors in the patient. Alternatively, the subject products may be used to bind to the immunoglobulins which bind to factor VIII proteins in the patient, so as to prevent the
35 immunoglobulins from diminishing the available factor VIII in the patient.

The hosts which are employed are microbial hosts, either prokaryotic or eukaryotic. Of particular interest for commercial fermentation are prokaryotic hosts of the Bacillus 5 genus, more particularly B. subtilis or B. licheniformis. Among eukaryotes of particular interest are yeasts, particularly of the genus Saccharomyces and Kluyveromyces, more particularly Kluyveromyces, and preferably K. lactis. Therefore, in accordance with this invention, the constructs 10 which are prepared and described will be exemplary for microbial organisms, but be directed to the preferred hosts, Bacillus and Kluyveromyces.

The particular hosts which are employed will desirably be industrial strains which are stable and have a 15 high production. The hosts may be modified to enhance yields. For example, with the Bacillus strains, problems with exoproteases or plasmid instability may be rectified. The exoprotease problem may be corrected by employing exoprotease deficient mutants (Kawamura and Doi, 1984). While these 20 authors modified B. subtilis, in the subject invention an improved strain of B. licheniformis has been produced (see Example 4). Where extrachromosomal maintenance is desired, opportunities for homologous recombination may be reduced by chromosomal deletion of sequences homologous to sequences 25 present in the plasmid. There may also be an interest in removing promoters or other regulatory sequences from the chromosome common to the plasmid.

The proteins of interest are those which aid in 30 correcting the clotting defect in providing a function lacking in the patient due to a defect in the endogenous factor VIII provision of the patient or due to endogenous antibodies which inactivate factor VIII (human), earlier defined as the combination of the 80 kDa and the 92 kDa proteins. The 80 kDa 35 subunit extends to the carboxy terminus from Glu-1649 through Tyr-2332 and the 92 kDa subunit extends from Ala-1 through

Arg-740, with a bridge between the two sequences having proteolytic cleavage sites. The bridge or central region referred to as the B-domain of factor VIII will range from Ser-741 through Arg-1648 (numbering according to Table 1).

5 The peptides of interest will include at least 10, more usually at least 12 amino acids. They may include the entire sequences of the 80 kDa and 92 kDa factor VIII subunits, and comprise fragments used to competitively bind antibodies to factor VIII, which will usually be homologous to
10 a factor VIII sequence. Otherwise, usually less than 10%, more usually less than 5% of the amino acid sequence will be modified from the naturally occurring sequence by mutations, that is conservative or non-conservative substitutions, deletions or insertions. The smaller the fragment, the fewer
15 the number of mutations which will be permissible, generally the total number of amino acids involved will be less than 15, more usually less than 10. A higher number of substitutions is permissible in cases where the substitutions are conservative, such as those amino acids which are aliphatic and which may be
20 divided into groups which are neutral, polar, acidic or basic and those amino acids which are aromatic.

The proteins which are prepared may be divided into the following categories:

25

The first category consists of fragments from about 8 to 60, more usually from about 12 to 50 amino acids which are able to compete with factor VIII for antibodies, so as to serve to remove from the patient antibodies which may bind
30 either to endogenous factor VIII or to factor VIII which is administered. These fragments may be used as fragments, or may be modified by joining them to various carriers, such as particles, other proteins, liposomes, or the like. The particular manner in which the fragments are modified will
35 depend upon the mode of administration and the particular purpose for which the fragments are used.

- The next group of proteins will be the subunits, which may be modified by truncation or extension, usually not exceeding 10% of the number of amino acids of the protein,
- 5 where the proteins will be active in a complex of the two subunits to provide for clotting activity. In the case the subunits are biological subunits and may be administered individually, it is intended that the subunit forms a complex with an endogenous subunit giving active factor VIII.
- 10 Alternatively, fused proteins may be prepared where all or a major portion (>90%) of a subunit may be fused to another protein, usually at least about 5 amino acids, particularly where the other protein may provide a signal sequence.
- 15 The next group of proteins will be proteins having sequences which individually are active in forming a clotting complex, but are connected, either head-to-head, head-to-tail, or tail-to-tail, either directly or by a bridge of fewer than about 900 amino acids, usually fewer than 500 amino acids,
- 20 more usually fewer than about 60 amino acids. The bridge may be a synthetic bridge, where a portion of the natural bridge is removed, so as to substantially shorten the spacing between the two subunits as compared to the natural precursor. Also, the subunits may be truncated at their N- and/or C-termini,
- 25 usually by not more than about 10 number %, more usually not more than about 5 number %. Desirably, the two subunits will be joined to each other or to the bridge by processing signals, which allow for enzymatic cleavage, particularly enzymatic cleavage in the intended patient, to provide for the
- 30 individual fragments. Either the naturally occurring processing signal or an alternative processing signal may be present, there being a large number of protease signals recognized by human hosts, e.g., K-R, R-R, etc.
- 35 The coding sequence may be obtained in a variety of ways. The sequence for the factor VIII precursor has been

reported in the literature (see also table I). Following the procedures described in the literature, either a genomic sequence or a cDNA sequence might be prepared of all or a portion of the factor VIII subunits. Thus, exons, fragments or
5 all of the cDNA for factor VIII precursor, synthetic sequences, or combinations thereof, may be employed for the encoding region. The sequences may be modified by in vitro mutagenesis, primer repair, resection, restriction, blunt ending, either by digestion or filling in, insertion of
10 adaptors, insertion into polylinkers, ligation with linkers, and the like. Thus, numerous techniques have evolved for modifying a coding sequence, to provide for deletions, insertions, and substitutions, such as transitions and transversions. The preparation of the coding sequence may be
15 done independently of the joining to the other members of the expression cassette or may be involved with the same process. Varying strategies may be employed, depending upon available sequences, vectors, the presence of restriction sites, or the like. Conveniently, the coding sequence may first be prepared,
20 followed by joining to the other members of the expression cassette.

At the 5' terminus of the coding sequence, a signal leader may be introduced. The signal leader will serve to provide for secretion of the desired protein into the nutrient
25 medium. The signal leader will normally be joined to the coding sequence by a processing signal, usually involving at least 2 codons, which code for a site which is recognized by an enzyme, which will cleave at that site. A large number of signal leaders have been studied in a variety of hosts. Signal
30 leaders which have been employed in the preferred Bacillus strains are alpha-amylases, exoproteases and levansucrase. Signal sequences which have been employed in the yeast strains, particularly the Kluyveromyces strains, include alpha-factor, either Saccharomyces or Kluyveromyces alpha-
35 factor, and synthetic leader sequences (see example 14).

The expression cassette will comprise a transcriptional and translational initiation region, joined to the 5' terminus of the coding sequence, so as to regulate transcription and translation of the coding sequence. Joined to the 3' end of the coding sequence will be a transcriptional and translational termination region, where the two regulatory regions will be functional in the intended expression host.

A number of transcriptional and translational initiation regions may be employed, but the regions will vary as to their efficiency in particular hosts and their effect on the growth characteristics of the particular host.

Transcriptional and translational initiation region for Bacillus of particular interest are alpha-amylases, exoproteases and levansucrase. Transcriptional and translational initiation regulatory regions for Kluyveromyces of particular interest are lactase, phosphoglycerate kinase, enolase etc. The termination regions are not as critical as the initiation regions, so that any convenient termination region may be employed. Thus, the termination region may be from the same gene as the transcriptional and translational initiation region or may be associated with a different gene, provided that the termination region is functional in the host. In many cases, the termination region may be associated with the gene from which the signal leader has been obtained or with a different gene.

The expression cassette may be used to transform the host directly or may be present as part of a vector. The vector is preferred since it provides for a number of conveniences. Most vectors will include one or more markers which allow for selection in the transformed host. Thus, one can select for those hosts which contain the expression construct as distinct from those hosts which have not been transformed. Markers for the most part provide for protection from biocides, particularly antibiotics, or impart prototrophy to an auxotrophic host. Illustrative markers include resistance to gentamycine (G418), kanamycin, tetracycline,

ampicillin, chloramphenicol, neomycin etc. The resistance genes will have appropriate transcriptional and translational regulatory regions to function in the expression host.

The vector will also have a replication system which is functional in the host, which may be low or high copy number. Depending upon the nature of the construct, the vector may be safely maintained as an extrachromosomal element or may become integrated into the host chromosome, particularly where the vector has regions of homology between DNA sequences of the vector and DNA sequences of the host. Thus, integration can be encouraged, by providing for a region of homology between the vector and the host and particularly where the replication system which is chosen is unstable and can only be maintained under selective pressure.

Conveniently, the vector may be a shuttle vector, where after each manipulation, the vector may be transferred to a cloning host, such as E. coli. In this manner, the constructions may be amplified and analyzed to ensure that the desired results have been obtained.

Once the expression vector has been completed, it may then be used for transformation. Various techniques for transformation exist for the desired species. Both Bacillus and Kluyveromyces may be transformed employing protoplasts in the presence of a fusogen, particularly polyethylene glycol. Alternatively the plasmids may be linearized and K. lactis transformed by them in a medium containing a transformation agent e.g. lithium acetate and a biocide e.g. G418. Cells may then be regenerated and selected for the presence of the expression cassette. The transformants may then be grown in an appropriate culture medium for production of the desired products.

The desired products may be isolated in a variety of ways. Where the product is retained in the cytoplasm, the cells may be harvested, lysed, and the protein extracted using

liquid-liquid extraction, chromatography, electrophoresis, or other conventional techniques. Where the product is secreted, the nutrient medium may be continuously circulated or partially withdrawn, and the desired product obtained as described with the lysed cells.

Once the product has been obtained into the desired purity, it may be brought in a pharmaceutical preparation adapted to its intended purpose. For use with a patient, it may be combined with a variety of physiologically acceptable additives, such as water, saline, phosphate buffered saline, or the like and administered to the patient, usually by injection, particularly intravascularly. The amount administered will vary widely depending upon the purpose of the protein, the status of the patient, the protein involved, its activity, as well as other conventional considerations. For the most part, the physiologically active subunits will be administered such that a level is attained of from about 0.1 to about 10 units per ml of blood.

Another aspect of the invention is the preparation of antibodies that recognize factor VIII subunit proteins. They may be prepared, using conventional techniques, by immunizing mammals with the proteins subject of the invention. Antibodies are obtained specific to the 80 kDa proteins and other to the 92 kDa proteins. When recovered from the blood they can be used in the production of hybridomas from which monoclonal antibodies can be produced. The antibodies of the present invention can be used in the diagnosis of factor VIII abnormalities, using any of the conventional competitive or direct assays on blood samples, which comprises bringing the sample into contact with the produced antibody and assaying for the presence or absence of a conjugate of the antibody with factor VIII antigen as a measure of the factor VIII protein in the sample.

The antibodies can also be used for the purification of factor VIII proteins which comprises bringing a liquid

phase comprising protein and a content of impurities into contact with a solid phase on which the produced antibody is immobilized, separating the liquid phase from the solid phase and eluting from the solid phase protein with a reduced 5 content of impurities.

The following examples are offered by way of illustration and not by way of limitation.

10

EXAMPLES

General Methods

1. Cloning techniques

15

For general cloning techniques reference may be made to the handbook of Maniatis et al., 1982.

Restriction enzymes are used as recommended by the manufacturer and are obtained either from New England 20 Biolabs (Biolabs), Bethesda Research Laboratories (BRL) or Boehringer Mannheim (Boehringer). In general 1 to 5 units of enzyme are needed to cleave 1 microgram of DNA.

For ligation, DNA fragments are run on an agarose gel, electroeluted, purified over a NACS column (BRL) as 25 recommended by the manufacturer, ethanol precipitated and incubated in a buffer containing 50mM Tris-HCl pH7.4; 10 mM MgCl₂; 10mM dithiothreitol; 1mM ATP and T4 DNA ligase (Biolabs).

For end-labeling, DNA cut with a restriction enzyme 30 is phenol-extracted and ethanol-precipitated. The phosphate is removed from the 5' end with Calf Intestine Alkaline Phosphatase (Boehringer) in 50mM Tris-HCl pH8.0 at 37°C. After inactivation of the enzyme at 65°C for 1 hr, followed by phenol extraction and ethanol precipitation, the DNA is 35 labeled using gamma-³²P-ATP (1-2 micromolar) in a buffer containing 50mM Tris-HCl, pH 7.6; 10mM MgCl₂; 5mM

dithiothreitol for 30 minutes at 37°C. Unincorporated nucleotides are removed by Sephadex G-50 chromatography.

Filling in of 5' overhanging restriction sites was done with the large fragment of DNA Polymerase I (Biolabs).

5 Incubation was for 15 minutes at 30°C in a buffer containing 10mM Tris-HCl pH7.7; 50mM NaCl; 10mM MgCl₂; 1mM dithiothreitol and unlabeled nucleotides at 20 micromolar.

Transformation of E. coli was either by the classical CaCl₂ technique (Maniatis et al., 1982) or exactly
10 as described by Hanahan (1983). Transformation of Bacillus or Kluyveromyces with foreign DNA was as described in the text.

2. Protein analysis. Sample preparation and electrophoresis

15 To 40 microliters of Kluyveromyces cell extract 10 microliters of sample buffer (0.313 M Tris-HCl pH6.8, 10% Sodium dodecylsulphate (SDS), 50% glycerol), 2.5 microliters of a saturated solution of Bromophenol Blue (BPB) in water and 2.5 microliters of beta-mercaptoethanol were added. The
20 samples were boiled for 2 min, centrifuged in an Eppendorf centrifuge for 2 min and subsequently layered on polyacrylamide gels according to Laemli (1970), consisting of a 7.5% separation gel and a 5% stacking gel. The gels were run at 60 mA until the BPB marker reached the bottom of the gel.

25 Samples of Bacillus cell extracts and extracellular proteins were dissolved in sample buffer (0.06M Tris-HCl pH6.8; 2% SDS; 10% glycerol). BPB and beta-mercaptoethanol were added as above. Samples were boiled for 5 min, centrifuged and subsequently layered and run on a 6%
30 polyacrylamide gel (5% stacking gel).

All gels depicted were run from top to bottom.

3. Protein blotting procedure

35 The proteins from the gel were blotted (Western blot) onto nitrocellulose filters (0.45 micrometers) using a

Biorad electroblot apparatus and a buffer containing 21 mM Tris; 160 mM glycine and 20% methanol, during 16 hrs at 150 mA. The blots were treated during 1 hr with a Bovine Serum Albumin (BSA)-solution (1% BSA in 50 mM Tris-HCl, pH7.5; 0.9% NaCl) and subsequently for 2.5 hrs with an antiserum in fresh BSA-solution. After 3 times rinsing with Tween 20 buffer (50 mM Tris-HCl, pH7.5; 0.9% NaCl and 0.05% Tween 20) the second antibody in fresh BSA-solution was added and incubated for 2 hrs followed again by 3 rinses with Tween 20 buffer.

10 The blots were stained depending on the enzyme coupled to the second antibody. In the case of horse-radish peroxidase staining was for 20 minutes with 0.6 mg/ml 4-chloro-naphthol and 0.015% H₂O₂ in a buffer containing 50 mM Tris-HCl, pH7.5 and 0.9% NaCl followed by rinsing in distilled water. When

15 alkaline phosphatase was used the blot was stained during a 15 min incubation with 0.033% nitro blue tetrazolium and 0.017% 5-bromo-4-chloro-3-indolyl-phosphate in 100 mM Tris-HCl, pH9.5; 100 mM NaCl; 5 mM MgCl₂.

20

4. Determination of chymosin

In the case of the 92 kDa-prochymosin fusion product part of the samples were treated at low pH in order to activate chymosin. To 500 microliters of cell extract 4M HCl

25 was added to lower the pH to approximately 2.0 and the sample was incubated at room temperature for 2 hrs. The pH was adjusted to 6.4 with 2.5M Tris. The samples were used for analyses on polyacrylamide gels and for determination of chymosin activity according to a standard milk clotting assay

30 (Foltman, 1970).

5. Preparation and characterization of monoclonal and polyclonal antibodies to factor VIII

35

Balb/c mice were immunized with purified human factor VIII-von Willebrand factor complex. Purification is

performed by an agarose gel filtration of cryoprecipitate or a factor VIII concentrate used for therapeutical purposes (Central Laboratory of The Netherlands Red Cross Blood Transfusion Service, Amsterdam, The Netherlands) as described 5 by Van Mourik and Mochtar (1970). Lymphocyte hybridization was performed as described by Galfré et al. (1977). Description of the techniques used for selection of clones producing monoclonal antibodies to factor VIII have been provided elsewhere (Stel et al., 1983).

10 Monoclonal antibodies to factor VIII were identified according to their reactivity with factor VIII polypeptides using a procedure adapted from Towbin et al. (1979). Batches as described above were first analyzed by SDS-polyacrylamide gel electrophoresis (Laemli, 1970). Then, separated proteins 15 were transferred electrophoretically (Western transfer) from the gel into a nitrocellulose sheet. The sheets were washed and incubated for 2 hrs with monoclonal antibodies and washed again. Then, the sheets were incubated with peroxidase-conjugated goat-antimouse immunoglobulins, washed and 20 incubated with 4-chloro-1-naphthol and H_2O_2 to detect antibody-bound peroxidase (see also above).

This procedure demonstrated that the following distinct monoclonal antibodies had been raised (Fig. 1). The antibody, designated CLB-CAG9, reacted with a polypeptide of 25 92 kDa and larger polypeptides. One antibody, CLB-CAG 65 reacted with the doublet of 79-80 kDa. The latter antibody did not react with the 92 kDa polypeptide and larger polypeptides. This demonstrates that the 79-80 kDa doublet possesses an epitope which is not present on the 92 kDa polypeptide.

30 In addition, monoclonal antibodies to factor VIII were identified according to their reactivity with factor VIII polypeptides, purified by immunoaffinity chromatography as previously described (Stel, 1984). The major portion of this preparation, devoid of von Willebrand factor, consists of the 35 79-80 kDa doublet. Traces of larger polypeptides are also present. Analysis of this preparation by immunoblotting

demonstrated that, besides monoclonal antibody CLB-CAg 9, also the antibodies CLB-CAg 21 and CLB-CAg 62 reacted with the 92 kDa and larger polypeptides.

To obtain polyclonal antibodies to factor VIII polypeptides, rabbits were immunized by standard procedures with the factor VIII preparation, purified by chromatography as described above. The antibodies thus obtained were identified by immunoblotting as described above using purified factor VIII-von Willebrand factor or purified factor VIII polypeptides, separated by SDS-polyacrylamide gel electrophoresis and subsequently transferred to nitrocellulose sheets as target proteins. This analysis demonstrated that three distinct polyclonal antisera had been raised (RH 63271, RH 63272 and RH 63275). These antisera reacted with the 79-80 kDa doublet, the 92 kDa polypeptide and larger polypeptides. It also shows that these antisera react with smaller polypeptides.

Example 1

20

SYNTHESIS OF A cDNA TO FACTOR VIII mRNA

Isolation of human liver polyA+ RNA:

25

During work with RNA care was taken to work Ribonuclease free. Sterile, disposable plasticware was used and all glassware was rendered nuclease free by heating overnight at 180°C. Wherever possible, solutions were incubated with 0.1% diethylpyrocarbonate for 30 min and

30 autoclaved.

Total RNA from 60 g of human liver tissue was prepared by the guanidium thiocyanate method as described by Chargwin et al. (1979). Homogenization of finely grounded frozen liver tissue in guanidium thiocyanate was done with a Polytron for 45 sec. The yield of total RNA was approximately 133 mg. The isolation of polyA+ RNA was done by a modification

of the procedure outlined by Aviv and Leder (1972). Total RNA was bound "batchwise" to an appropriate amount of oligo(dT)-cellulose (Collaborative Research, type T2) in 50 mM Tris-HCl (pH 7.5); 1 mM EDTA; 0.2 % SDS; 0.4 M NaCl. A column was prepared from this preparation and polyA⁺ RNA was eluted with the same buffer lacking NaCl. A second cycle of absorption of polyA⁺ RNA to oligo(dT)-cellulose was performed using less stringent conditions (buffer containing 0.25 M NaCl), in order to separate polyA⁺ RNA from remaining ribosomal RNA. The final yield of polyA⁺ RNA was about 830 micrograms. Analysis of this preparation by agarose gel electrophoresis showed that it did not contain ribosomal RNA.

Primer-directed cDNA synthesis:

15

Synthesis of factor VIII cDNA was based on the method of Gubler and Hoffman (1983) as modified by Toole et al. (1984). Specific primers for cDNA synthesis were synthesized with a Biosearch DNA synthesizer (Figure 2). Human liver polyA⁺ RNA (20 micrograms in distilled water) was mixed with a 10-fold molar excess of single stranded primer in 10mM CH₃HgOH in a total volume of 35 microliters and incubated for 10 min at room temperature. Beta-mercaptoethanol was added to a final concentration of 47 mM and then RNasin (Promega-Biotec) to 100 U/ml (total volume 47 microliters). Then 50 microliters of 2xRT buffer (0.1M Tris-HCl pH8.3 (43°C); 10mM MgCl₂; 150mM KCl; 60mM beta-mercaptoethanol; 1mM each of dATP, dGTP and dTTP and 0.2mM of dCTP) and 0.4 microliters of water were added. After 2 min incubation at 50°C followed by 2 min at 43°C, 2.6 microliters of super RT (Anglian Biotechnology, 17 U/microliter) were added. First strand synthesis was for 45 min at 43°C. The reaction was stopped on ice and 300 microliters of a solution containing 40mM Tris-HCl (pH7.5); 10mM MgCl₂; 75mM KCl and 25 micrograms/ml BSA (BRL, Desoxyribonuclease and Ribonuclease free) were added. For

second strand labeling 2 microliters of alpha-³²P-dCTP (3000Ci/mmol; 10mCi/ml) were added, followed by 8 microliters of DNA Polymerase I (10 U/microliter; Biolabs) and 4 microliters of RNase H (10 U/microliter; Anglian). After 60 min at 11°C, 4 microliters of 20mM ATP and 1 microliter of T4 DNA ligase (400 U/microliter; Biolabs) were added and incubation continued for 120 min at 18°C. The reaction was stopped by the addition of 30 microliters of 0.5M EDTA and 4 microliters of 10% SDS. The mixture was extracted with phenol and the DNA precipitated with isopropanol. Unincorporated nucleotides were removed on a Sephadex G-50 column. The final yield was 400 ng of double stranded cDNA.

Isolation of partial factor VIII cDNA clones:

15 The double stranded cDNA was digested with a combination of two restriction enzymes as indicated in Figure 2. After digestion, cDNA fragments were separated according to size on a Sepharose CL-4B column. Samples of fractions from the column were run on an agarose gel and the length estimated by autoradiography. Material larger than 600 bp was used for subsequent cloning experiments. Fragments A and B were ligated into pAT153 (Twigg and Sherrat, 1980) C and D in pACYC177 (Chang and Cohen, 1978) and E in pUC19 (Yanish-Perron et al., 1985). Before ligation to cDNA fragments, all plasmids were digested with the corresponding two restriction enzymes and the vector fragment isolated by agarose gel electrophoresis, followed by electroelution and DEAE-Sephacel chromatography.

 After ligation, E. coli DH1 was transformed exactly as described by Hanahan (1983). About 100 colonies with plasmids containing an insert were obtained per nanogram of cDNA. Colonies were transferred to nitrocellulose filters (Grunstein and Hogness, 1975), lysed and hybridized to radioactively labeled probes (Figure 2). Clones which were derived from cDNA synthesis with fragment 1 as primer were analysed with fragment 2. Clones synthesized with fragment 2

were analysed with fragment 3, etc. Labeling of fragments was with gamma-³²P-ATP and polynucleotide kinase (Maniatis et al., 1982). Prehybridization of filters was at 20°C in a solution containing 6 x SSC, 5 x Denhardt's (0.1% Ficoll; 0.1% polyvinyl pyrrolidone; 0.1% BSA) and 100 micrograms of E. coli DNA per ml. Hybridization was in the same solution containing 10 ng of labeled fragment per ml overnight at 20°C. Filters were washed in 6 x SSC at 37°C and autoradiographed. About 1 in 1000 colonies was positive and correct clones were isolated for all cDNA subfragments. Clones from fragments A, B, C, D and E (Figure 2) were called pCLB81, pCLB82, pCLB83, pCLB84 and pCLB85, respectively.

Construction of a full length factor VIII cDNA:

15

Clone pCLB85 is missing a small DNA fragment from the factor VIII-encoding sequence from the ATG start codon up to the SacI site at position 16 (Table I). The missing nucleotides were provided by chemical synthesis of a DNA fragment thereby creating a SalI site upstream of the ATG codon such that it reads 5'GTCGACATGCAAATAGAGCTC3'. To this end pBR322 was cleaved with AvaI and the AvaI site was filled in with the large fragment of DNA Polymerase I. After a second cleavage with PvuII the remaining pBR322 fragment was circularized giving plasmid PI. The 1.9kb factor VIII cDNA-specific BamHI-SacI fragment from pCLB85 was ligated to PI cleaved with BamHI and BanII. After ligation a chemically synthesized DNA fragment (CGGTCGACATGCAAATAGAGCT) was added, and the mixture was incubated with the large fragment of DNA polymerase I and all four desoxyribonucleotides. After phenol extraction, followed by digestion with SalI and another phenol extraction the fragment was ligated. The resulting plasmid pCLB86 now contains a SalI site in front of the factor VIII ATG start codon.

35

Plasmids pCLB81, 82, 83, 84 and 86 are now assembled to a full length factor VIII clone in 3 steps. First, the

inserts of pCLB84 and 86 are combined in pUC19 cleaved with SalI and PstI (pCLB87). Then, plasmids pCLB81 and 82 are cleaved with EcoRI plus HpaI and BamHI plus EcoRI, respectively, and ligated into pEP121 (Enger-Valk, 1984) giving pCLB88. Then, the inserts of pCLB83, 87 and 88 are combined to a cDNA clone containing the complete protein-encoding part of the factor VIII gene as a SalI-HpaI fragment in pEP121 (pCLB89).

The factor VIII cDNA insert in pCLB89 was sequenced according to the methods of Sanger et al. (1977) and Maxam and Gilbert (1980). The sequence is shown in Table I.

Example 2

15 CONSTRUCTION OF A BACILLUS SECRETION VECTOR WITH THE 20 BACILLUS LICHENIFORMIS ALPHA AMYLASE GENE EXPRESSION CASSETTE

Cloning of the alpha-amylase gene from
20 B. licheniformis strain T5 in the pUB110 (Matsumura et al., 1984) vector yielded the recombinant plasmids pGB33 and pGB34 as described in European patent application EP-A-0134048.

The inserted B. licheniformis restriction fragments from pGB33 to be sequenced were subcloned into the
25 multicloning site of M13mp8 and M13mp9 (Messing and Vieira, 1982) and nucleotide sequences were determined by the di-deoxy chain termination method of Sanger et al (1977).

The promoter, Shine-Dalgarno sequence and the signal sequence of the cloned alpha-amylase gene are described in
30 European patent application EP-A-0224294.

In order to place a gene for bovine chymosin behind this alpha-amylase transcriptional and translational initiation regulating sequences we digested pGB34 and pUR1523, an E. coli plasmid that harbours the gene for bovine chymosin
35 (see European patent application EP-A-0077109), with PstI. After ligation and transformation to competent B. subtilis 1A-

40 (BGSC, Ohio U.S.A.) cells, plasmid DNA was isolated from selected transformants. After restriction endonuclease analyses the correct recombinant plasmid called pLC28 was obtained. Cutting pLC28 with the restriction endonuclease
5 ClaI, followed by resection with the exonuclease Bal-31, ligation and transformation yielded pLC83 (see European patent application EP-A-0134048).

From recent literature (Ulmanen et al. (1985),
10 Shiroza et al. (1985)) it was known that the level of secretion and production of a heterologous gene fused to an alpha-amylase based secretion vector is also depending on the codon position of the mature alpha-amylase at which the heterologous gene is joined. Therefore, we incubated a M13mp8
15 alpha-amylase harbouring clone (called pPB4) with SstII followed by resection with the exonuclease Bal-31, cutting with SmaI, ligation and transformation to competent E. coli JM101 cells (Yanisch-Perron et al., 1985). This procedure yielded a so-called alpha-amylase "leaderbank". This means a
20 set of M13mp8 derived clones where the number of amino acids (called n) behind the maturation site of the alpha-amylase protein varies.

The M13mp8 "leader" clones with the correct reading frame have been used to place the (pro)chymosin gene in phase
25 behind the alpha-amylase transcriptional, translational and secretion regulatory sequences. These fusion plasmids were called pLCn for the chymosin constructs and pLPn for the prochymosin constructs, respectively, where n stands for the number of alpha-amylase amino acids behind the maturation
30 site.

The pLCn and pLPn constructs were modified in such a way, that the EcoRI site, originating from plasmid pGB33 was removed by exonuclease Bal-31 action after cutting with EcoRI.
Furthermore, the pLC constructs have been modified in such a
35 way that they contain unique restriction endonuclease sites directly behind the "leader" sequences, and a recognition site

for signal peptidase directly behind the "leader" sequences. To fulfill these two requirements four oligodeoxyribonucleotides, I to IV, were chemically synthesized:

- 5
a. I : 5'-AATTCGCGGCCGCC-3' (no. 104 1526 GBR 22774)
II : 5'-CGGGTCGACTCAAGCTTA-3' (no. 104 1527 GBR 22775)
III : 3'-GCGCCGGCGGGCCAGCT-5' (no. 104 1524 GBR 22773)
IV : 3'-GAGTTCGAATTTAA-5 (no. 104 1523 GBR 22772)
10 b. EcoRI XmaIII XmaI/SmaI SalI HindIII

5' AA TTC GCG GCC GCC CGG GTC GAC TCA AGC TTA 3'

← I → ← II →

15

3' G CGC CGG CGG GCC CAG CTG AGT TCG AAT TTA A 5'

← III → ← IV →

20 glu phe ala ala ala arg val asp ser ser leu asn ser

These fragments constitute the multicloning site and they were joined to pLC constructs yielding the secretion vectors of the pOL5-n type (Fig. 3). This plasmid serves as a vehicle for
25 subsequent introduction (see Example 3) of expressible genes encoding proteins with factor VIII activity, the plasmid being suitable for transforming Bacillus. n in the plasmid designation refers to the number of amino acids from the mature alpha amylase present behind the signal sequence, Fig.
30 3 illustrating a plasmid that will produce 21 amino acids in this section. The number of amino acids in this section does not affect the ability of the plasmid, when carrying a factor VIII protein gene to express factor VIII protein in a Bacillus transformant. Bacillus subtilis was transformed with pOL5-21
35 and the transformant deposited on 14 July 1986 in the Centraal Bureau voor Schimmelcultures (the CBS), at Oosterstraat 1,

3742 SK BAARN, NETHERLANDS where it has been given the deposit number CBS-304.86.

By using the pOL5-n expression vector having 5 sequences homologous with the host chromosome of B. licheniformis T5 strain we observed both segregational and structural instability of the plasmids.

To circumvent homologous recombination between the alpha-amylase sequences on the B. licheniformis host cell chromosome on one hand and the same sequences on the expression vector we have decided to delete these sequences.

Firstly we constructed a Bacillus licheniformis host strain which lacks the alpha-amylase promoter and the complete alpha-amylase chromosomal gene (see Example 4).

Secondly we removed from the expression vector the remaining homologous sequences upstream the alpha-amylase promoter. Therefore the pOL5-n vectors were digested by EcoRV and BglI, ligated and transformed to protoplasts of Bacillus subtilis 1A40 (BGSC, Ohio U.S.A.). Plasmid DNA was isolated from neomycin resistant transformants. After restriction endonuclease analyses the right recombinant plasmids designated pOL5-delta were obtained (see Figure 4).

Protoplasts of Bacillus licheniformis T9 strain (see Example 4) transformed with these pOL5-delta vectors yielded neomycin resistant transformants. These T9 transformants were subjected to a plasmid stability test as described by Andreoli, 1985.

The plasmid DNA population was analysed in more detail by restriction enzyme mapping. After about 50 generations less than 5% of the pOL5-delta vectors have been cured in contrast to T9 cells harbouring pOL5 plasmids where up to 90% of the plasmids have been cured.

These results indicate that the pOL5-delta expression vectors are stable in the genetic background of the T9 Bacillus licheniformis host.

Example 3

INTRODUCTION OF THE HUMAN FACTOR VIII cDNA INTO A BACILLUS
VECTOR AND ANALYSIS OF EXPRESSION OF THE 92 kDa FACTOR VIII
5 SUBUNIT IN BACILLUS SUBTILIS.

In pOL5-21 the 92 kDa subunit gene is inserted
yielding pOL92 (see Figure 5A,B). After transformation of
protoplasts from Bacillus subtilis (Chang and Cohen, 1979)
10 followed by colony hybridization analysis (Grunstein and
Hogness, 1975) and plasmid restriction enzyme mapping, we
obtained the constructs pOL92-T-21 and pOL92-A-21,
respectively.

To examine the expression of factor VIII
15 polypeptides, Bacillus subtilis transformants were grown in an
appropriate medium e.g. Trypton soya broth at 37-40°C with
aeration in the presence of neomycin (10 micrograms/ml).
Samples of culture were centrifuged and lysed as follows:
After centrifugation of a 2 ml culture, cells were resuspended
20 in 0.1 ml of a buffer containing 50mM Tris-HCl, pH7.5 and 50mM
EDTA. One microliter of 100mM PMSF was added and egg white
lysozyme to a final concentration of 2mg/ml. After 5 min at
37°C another microliter of 100mM PMSF was added and incubation
continued for 5 min at 37°C. The reaction was terminated by
25 the addition of sample buffer.

The SDS-polyacrylamide gel analysis of these samples
is shown in Fig. 6. When analyzed by SDS-polyacrylamide gel
the cell lysate from pOL92-T-21 contained bands of molecular
weights of 46 kDa, 58 kDa and 88 kDa, respectively, which are
30 much stronger than in the control. The molecular weight of
the largest pOL92-T-21 encoded protein is 88 kDa. Based on the
amino acid composition the large subunit polypeptide has a
molecular weight of 85 kDa (Vehar et al., 1985), while for the
human fragmentation product a molecular weight of 92 kDa was
35 measured. Differences in molecular weight might be accounted

for partly by experimental error, by differences in protein modifying reactions like glycosylation, and/or differences in processing of the translated protein products. Thus, this result demonstrates that we have produced large subunit factor 5 VIII protein in Bacillus.

Bacillus subtilis transformed with pOL92-T-21 was deposited in the CBS on 14 July 1986 where it was given the deposit number CBS-303.86.

10

Example 4

CONSTRUCTION OF AN EXOPROTEASE DEFICIENT AND ALPHA-AMYLASE
DEFICIENT BACILLUS LICHENIFORMIS HOST

15 Bacillus licheniformis produces at least two major extracellular proteases, an alkaline protease (or subtilisin) and a neutral protease. Therefore strain T5 (European patent application EP-A-0134048) was mutagenized by two subsequent cycles with ultraviolet radiation according to the method of 20 Miller (1972) and plated on Spizizen glucose minimal medium supplemented with 0.4% casein. Seven of 12.000 colonies screened produced a smaller halo after the first cycle. These seven mutants were subjected to the second cycle of UV mutagenesis. Two of 10.000 colonies screened showed no halo or 25 extracellular protease activity on the 0.4% casein plates. These double mutants were characterized for both total exoprotease production in liquid cultures as described by Kawamura and Doi (1984) as well as the alpha-amylase production by measuring the production of reducing sugars from 30 starch (Ghandi and Kjaergaard, 1975) to exclude the isolation of mutants defective in secretion. One such isolate, which was alpha-amylase proficient and of which the exoprotease activity was beyond detection was designated strain T6.

The next step was to construct a Bacillus
35 licheniformis T6 derivative which lacks the alpha-amylase

promoter and the complete alpha-amylase structural gene. To create such deletion mutation (designated delta Amy) the recombinant plasmid pGB33 as described in European patent application EP-A-0134048 was digested with Hind III and BglI, 5 ligated and transformed to competent cells from Bacillus subtilis strain 1A40 (BGSC, Ohio, U.S.A.). Plasmid DNA was isolated from neomycin resistant alpha-amylase negative transformants. After restriction endonuclease analyses the correct recombinant delta Amy plasmid called pGB33delta was 10 obtained. This plasmid contains portions of the 5' and 3' flanking sequences of the Bacillus licheniformis alpha-amylase gene. To replace the wild-type alpha-amylase gene with the delta Amy deletion mutation the pGB33delta plasmid was transformed into Bacillus licheniformis T6 protoplasts as 15 described in Example 5. Two of 20.000 neomycin resistant transformants showed no extracellular alpha-amylase activity and retained the pGB33delta plasmid, indicating that the frequency of gene conversion was very low. The two alpha-amylase deficient Nm^R transformants were cured of the 20 plasmid by growing for 20 generations in minimal medium as described by Meyer and Fiechter (1985) in the absence of antibiotic selection.

Neomycin sensitive cells cured of the plasmid appeared at a frequency of about 2%. One of these colonies was 25 designated strain T9. We used Southern blotting and hybridization (Maniatis et al. 1982) to confirm that the T9 strain carried the 1.9 kb deletion of the alpha-amylase gene. The construction of the alpha-amylase deficient Bacillus licheniformis T9 host is presented in Figure 7.

30 An additional advantage of this T9 strain is when this host strain harbours a Bacillus licheniformis alpha-amylase secretion vector e.g. pOL5 (see Example 2) there exists no longer competition for binding RNA polymerase molecules between the chromosomal alpha-amylase promoter and 35 the alpha-amylase promoter located on the plasmid DNA.

Example 5

PROTOPLAST TRANSFORMATION AND REGENERATION OF BACILLUS LICHENIFORMIS STRAINS.

5

Bacillus licheniformis T5 is an industrial strain used for the production of extracellular enzymes, which has been described in European Patent application EP-A-0134048.

Bacillus licheniformis strain T5 was grown overnight
10 in 50 ml NBSG-X medium (Thorne and Stull, 1966) at 37°C. The culture was diluted 1:1 with fresh NBSG-X medium and grown for another 1-1.5 hrs. After centrifuging for 10 min at 5000 rpm in a Sorvall type GSA rotor and resuspending in 10 ml of SMM buffer, containing 1.5M sucrose; 0.06M MgCl₂ and 0.06M
15 maleate, the cells were protoplasted by incubating for 2 hrs at 37°C in the presence of 2 mg/ml of lysozyme.

The protoplasts were spun down (10 min at 5000 rpm), resuspended in 5 ml of SMML buffer (L-broth in which 1.5M sucrose; 0.06M MgCl₂ and 0.06M maleate has been dissolved),
20 mixed and repelleted. After being resuspended, the protoplasts were incubated for 2 min in 30% polyethylene glycol 6000 with 1 microgram of plasmid pGB33 (see Example 2).

After this incubation a 1:3 dilution with SMML medium was carried out and, after centrifugation, the
25 protoplast-containing pellet was resuspended in 1 ml of SMML medium. Then, 0.1 ml aliquots were plated on regeneration agar plates containing 0.7% (w/v) K₂HPO₄; 0.3% (w/v) KH₂PO₄; 0.125% (w/v) (NH₄)₂SO₄; 0.035% (w/v) MgSO₄·7H₂O; 1.5% (w/v) agar; 3.7% (w/v) KCl; 0.1% (w/v) glucose; 0.01% (w/v) BSA
30 supplemented with 0.1% (w/v) spore solution containing 0.2% (w/v) MnSO₄; 0.2% (w/v) ZnSO₄; 0.2% (w/v) CoCl₂; 0.5% (w/v) FeSO₄; 6% (w/v) NaCl and 0.5% (w/v) CaCl₂. Moreover, these plates contained 10 micrograms/ml of neomycin.

After incubation at 37°C for at least 72 hrs, the
35 plates were replica plated on heart-infusion agar plates

containing 10 micrograms/ml of neomycin. Transformants were analysed on their DNA content by isolation of DNA as described by Holmes and Quigley (1981) followed by characterisation on agarose gels. Among the transformants 5% were found not to contain any plasmid DNA, but they contained the complete pGB33 DNA as an integral part of their genome. Plasmid pGB33 was found to have been integrated into the chromosome by homologous recombination between the alpha-amylase sequences on the B. licheniformis T5 chromosome on one hand and the same sequences on the plasmid pGB33 on the other hand.

Example 6

15 CONSTRUCTION OF A BACILLUS INDUSTRIAL STRAIN SUITED FOR EFFICIENT CHROMOSOMAL INTEGRATION OF HETEROLOGOUS GENES.

The plasmid pBC16 is derived from Bacillus cereus and is highly homologous to pUB110 as described by Polak and Novick (1982). The main difference between pBC16 and pUB110 is that pBC16 contains a tetracyclin resistance gene where pUB110 has a neomycin resistance gene.

B. licheniformis T5 transformants, as described in Example 5, harbouring two amylase genes and one copy of pUB110 in their chromosome, were transformed using pBC16 linearized with XbaI, according to the protoplast transformation protocol as described in Example 5. 10 micrograms/ml of tetracyclin were included in the regeneration plates. Selection for tetracyclin resistant colonies resulted in a strain where the chromosomally located neomycin gene derived from pUB110 was replaced by the tetracyclin gene of pBC16 by double reciprocal recombination.

The resulting strain T5-16 can be used as a suitable strain for efficient chromosomal insertion of pUB110-derived plasmid constructs. Strain T5-16 is tetracyclin resistant and neomycin sensitive, and has approximately 3kb of DNA sequences

homologous to PUB110 in its genome.

To obtain an exoprotease deficient B. licheniformis strain suited for chromosomal integration of heterologous genes the strain T5-16 and strain T6 (see Example 4) were protoplasted and fused as described in European patent application EP-A-0134048.

The obtained tetracyclin resistant fusants were screened for exoprotease activities and characterized by genomic analysis using ³²P-labeled alpha-amylase encoding DNA probe. A fusant, which is tetracyclin resistant, exoprotease negative and has approximately 3kb of DNA sequences homologous to PUB110 in its genome, is designated strain T6-16.

15

Example 7

INTEGRATION OF THE HUMAN FACTOR VIII GENE INTO THE BACILLUS LICHENIFORMIS CHROMOSOME; EXPRESSION AND SECRETION OF 92 kDa FACTOR VIII SUBUNIT.

20

Bacillus licheniformis strain T5-16 was transformed with the plasmid pOL92-T-21 and pOL92-A-21 according to the protocol as described in Example 5. Transformants were selected on regeneration plates containing 5 micrograms/ml of neomycin. None of the transformants was found to contain plasmid DNA, in all cases integration of plasmid DNA by homologous recombination into the genome of strain T5-16 occurred.

To examine the expression of factor VIII polypeptides these T5-16 transformants were grown in heart-infusion medium, supplemented with 0.5% of starch. After 3 days of incubation at 40°C samples for analysis on protein gels were prepared as follows: 3 ml of fermentation broth was centrifuged and from the supernatant fraction protein was precipitated by addition of 5% trichloroacetic acid (TCA); the precipitated material was resuspended in 1% TCA and pelleted again by centrifugation. The protein from 25 ml of supernatant

was then dissolved in 0.5 ml of sample buffer. The pellet fraction of 6 ml of culture broth was dissolved in 0.7 ml of sample buffer. The composition of both supernatant and cell pellet fraction was studied by analysing 60 and 50 microliters, respectively, on 6% SDS-polyacrylamide gel as shown in Figure 8.

From the result it can be concluded that Bacillus licheniformis can synthesize and secrete factor VIII 92 kDa proteins as intact products because a protein band is clearly visible in cell pellet and supernatant protein fractions of B. licheniformis T5-16 cells transformed with pOL92-A-21 and pOL92-T-21, while in the control (strain T5-16) these bands are not seen.

Example 8

15

SYNTHESIS AND SECRETION OF THE 80 kDa AND 92 kDa SUBUNIT PROTEINS USING THE BACILLUS LICHENIFORMIS ALPHA-AMYLASE SECRETION VECTOR pOL5-DELTA

20

To create a Bacillus expression plasmid containing the 80 kDa subunit protein the following three DNA fragments were combined:

(i) the 4.6 kb SalI-BalI fragment from pOL5-40 delta containing the alpha-amylase promoter, leader, terminator, the neomycin resistance gene and a Bacillus origin of replication.

(ii) a 0.6 kb SalI-HincII fragment from a M13mpl8-80 kDa clone containing the N terminal part of the 80 kDa subunit preceded by an ATG codon and a SalI restriction site as described in Example 13.

(iii) a 2.3 kb HincII-EcoRV fragment from pGB852 containing most of the 80 kDa coding sequence and the lactase terminator.

After ligation of these three fragments the recombined DNA molecules are introduced in Bacillus by means of protoplast transformation.

Plasmid DNA was isolated from neomycin resistant transformants as described by Andreoli (1985) followed by restriction endonuclease mapping. The plasmid searched for was called pOL80-T-40delta and its structure is shown in Figure 9.

To examine the expression of factor VIII 80 kDa protein the obtained Bacillus subtilis 1A40 transformants were grown in Trypton soya broth medium with aeration in the presence of neomycin (10 micrograms/ml).

- 10 After overnight cultivation at 37°C protein samples were prepared and analysed as described in the General Methods section. Figure 10C shows the detection of the 80 kDa protein in the cell lysates of pOL80delta B. subtilis transformants in contrast to the pOL5delta control strains.
- 15 This result demonstrates that we have produced factor VIII 80 kDa protein in Bacillus subtilis.

To remove the homologous DNA sequences between the Bacillus licheniformis chromosome of the T9 strain and the plasmid pOL92-T-21 we followed the same procedure as described in Example 2 for the construction of the expression vector pOL5delta. The plasmid thus obtained, pOL92-T-21delta, has a structure as shown in Figure 5C.

- Also the 7.4 kb SalI-HpaI fragment from pCLB89 containing the full length factor VIII cDNA and the 5.8 kb
- 25 SalI-EcoRV fragment from pGB861 containing a fusion between the 92 kDa and 80 kDa proteins (see Example 14) were introduced in the alpha-amylase secretion vector pOL5delta in a similar way as described in Example 3. The plasmids thus obtained were called pOLF3-2delta and pOL92/80-21delta,
- 30 respectively.

To examine the expression of factor VIII polypeptides the Bacillus T9 transformants harbouring pOL5-21delta, pOL80-T-40delta and pOL92-T-21delta, respectively, were grown in double strength minimal medium (Anagnostopoulos and Spizizen, 1961) supplemented with 0.5% of yeast extract. After two days of cultivation at 40°C protein samples were

prepared for analysis on SDS-polyacrylamide gel, blotted onto a nitrocellulose filter and incubated with appropriate antibodies as described in the General Methods section. From the results shown in Figure 10A and 10B it can be concluded
5 that transformed Bacillus licheniformis T9 can synthesize and secrete factor VIII specific 92 kDa and 80 kDa polypeptides, respectively.

Example 9

10

EXPRESSING A FUSION PROTEIN BETWEEN THE 92 kDa AND 80 kDa
FACTOR VIII SUBUNITS USING A BACILLUS PLASMID CONSTRUCT

After fermentation of Bacillus transformants harbouring
15 respectively pOLF8-2delta and pOL92/80-2Idelta (see Example 8) only traces of factor VIII specific polypeptides could be detected.

The factor VIII DNA encoding a 92/80 kDa fusion protein was put behind the strong Bacillus licheniformis 3-4
20 promoter (see EP-A-0224294) as follows:

The 5.6 kb Sall-NsiI fragment encoding the 92/80 kDa fusion protein from pGB861, the 3.9 kb XbaI-XmaIII fragment encoding the 3-4 promoter, the neomycin resistance gene and an oligonucleotide with the sequence 5'dGGCCTGCA were ligated and
25 transformed to protoplasts of Bacillus. This transformation yielded the plasmid pPROM92/80PB (see Figure 11).

Bacillus strains harbouring pPROM92/80PB were cultured for at least 2 days at 40°C as described in Example 8 and protein samples were prepared and analysed for
30 expression of factor VIII polypeptides as described in the General Methods section.

Figure 10D shows the detection of the 92/80 kDa fusion protein in the cell lysate of pPROM92/80PB Bacillus transformants in contrast to the Bacillus control strains.

35

Example 10

CONSTRUCTION OF PLASMID pGBdLAC CONTAINING THE LACTASE
PROMOTER AND TERMINATOR SEQUENCES.

5

Chromosomal DNA was isolated from Kluyveromyces
lactis strain CBS 2360 (Das and Hollenberg, 1982), cleaved
with XhoI, and separated according to size on a sucrose
gradient. Fractions containing the lactase gene were detected
10 with a LAC4 probe (Breunig et al., 1984) after spotting the
DNA on a nitrocellulose filter. The lactase gene containing
DNA was cloned into the XhoI site of plasmid pPA153-215
(Andreoli, 1985) giving rise to plasmid pPA31. An XbaI
fragment of pPA31 containing the lactase gene was subcloned in
15 the XbaI site of pUC19 (Yanisch-Perron et al., 1985) which
yields plasmid pUCla56. The EcoRI-XbaI fragment containing the
3' end of the lactase gene (Breunig et al., 1984) was cloned
as a separate fragment in pUC19. In the EcoRI-XbaI fragment a
ClaI site is present which is normally methylated in a dam +
20 strain and is about 100 nucleotides downstream of the EcoRI
site. To be able to cut at this ClaI site, the plasmid was
transfected to a dam(-) E. coli strain, e.g. strain GM113
(Arraj and Marinus, 1983; Marinus and Morris, 1973). Plasmid
DNA was obtained and the ClaI-XbaI fragment containing the 3'
25 end of the gene isolated. In addition, the XbaI-ClaI fragment
containing the 5' end of the lactase gene was isolated from
pUCla56. Both fragments were simultaneously ligated into the
XbaI site of pUC19, giving rise to plasmid pGBdLAC (Fig. 12).

30

Example 11

CONSTRUCTION OF A K. LACTIS EXPRESSION VECTOR CONTAINING THE
92 kDa FACTOR VIII SUBUNIT.

35

As a first step in the synthesis of a 92 kDa subunit
expression plasmid the 1.4kb Hind III-PstI fragment,
containing the 3' end of the lactase gene (Breunig et al.,
1984), was ligated onto the 0.4kb BamHI-Hind III fragment

(nucleotides 1875 to 2283 as in Table I) of factor VIII cDNA and ligated into plasmid pSP65 (Melton et al., 1984) cleaved with BamHI and PstI, giving rise to plasmid pGB840. Next, a stopcodon which is in phase with the factor VIII reading frame 5 was introduced into the Hind III site. To this end an oligonucleotide with the sequence 5' AGCTGAGGGCCCTC was synthesized which contains in addition to the TGA stopcodon an ApaI recognition site (GGGCCC). It was ligated into the Hind III site of pGB840 yielding plasmid pGB841. From the latter 10 plasmid the 1.2kb BamHI-EcoRV fragment was isolated and ligated into the 7.4kb BamHI-EcoRV fragment of plasmid pCLB86. This gives rise to plasmid pGB842 containing the complete 92 kDa subunit encoding DNA containing a stopcodon at the Hind III site in pAT153. To create a K. lactis expression plasmid 15 containing the 92 kDa subunit protein the following 3 fragments were combined:

- (i) the 3kb SalI-EcoRV fragment from pGB842 containing the 92 kDa-subunit DNA and the lactase terminator.
 - (ii) A 0.25kb SnaBI-SalI fragment containing the part of the 20 lactase promoter from position - 282 (Breunig et al., 1984) to position -26 at which a SalI linker was introduced by the method described before (EP-A-0096430).
 - (iii) A 8.3kb SnaBI-EcoRV fragment from a derivative of pGBdLAC containing a G418 resistance gene (see EP-A-0096430).
- 25 The resulting plasmid was called pGB850 (Fig. 13).

E. coli transformed with pGB850 was deposited in the CBS on 14 July 1986 where it was given the deposit number CBS-305.86.

Example 12

30

CONSTRUCTION OF A K. LACTIS EXPRESSION VECTOR CONTAINING A FUSION BETWEEN THE 92 kDa-SUBUNIT AND PROCHYMOSIN.

The prochymosin cDNA was put in the same reading 35 frame behind the BamHI site at position 1875 of the factor VIII cDNA (Table I) as follows: The 11.0 kb EcoRV-BamHI

fragment of pGB850 was isolated and the BamHI site filled in using the Klenow fragment of DNA polymerase I and all four deoxynucleotide triphosphates. Next, the 2.3kb Sall-HindIII fragment containing the prochymosin gene from plasmid pGB131 5 (EP-A-0096430) was isolated and the Sall and Hind III sites filled in as above. Both fragments were ligated yielding plasmid pGB851 which contains the prochymosin-encoding DNA downstream of the 92 kDa-encoding DNA (Fig. 14).

10

Example 13

CONSTRUCTION OF K. LACTIS EXPRESSION VECTORS FOR THE 80 kDa SUBUNIT PROTEIN.

15

A vector was constructed for the expression of a lactase-80 kDa fusion protein. Two fragments were used for the 80 kDa expression plasmid:

- (i) A derivative of plasmid pGBdLAC containing the G418 resistance gene (above) was linearized with ClaI.
- 20 (ii) Factor VIII plasmid pCLB88 was cleaved with BamHI and HpaI and the 2.7kb fragment from position 4749 to position 7439 in the factor VIII cDNA sequence (Table I) was isolated. Both fragments were made blunt using the Klenow fragment of DNA polymerase I and all four deoxyribonucleotides, ligated 25 and E. coli HB101 transformed. Plasmids containing the 80 kDa-encoding part in the right orientation and in frame behind the lactase-encoding sequences were identified by restriction enzyme digestions and DNA sequence analysis according to Maxam and Gilbert (1980). The resulting plasmid pGB852 is shown in 30 Fig. 15.

E. coli transformed with pGB852 was deposited in the CBS on 14 July 1986 where it was given the deposit number CBS-306.86.

In another expression vector an ATG codon preceded by a Sall 35 restriction site was introduced in front of the protease cleavage site of the 80 kDa subunit protein:

a BamHI-HindII fragment from position 4749 to position 5584 of the factor VIII cDNA sequence was introduced in M13mpl8 (Messing and Vieira, 1982) cleaved with BamHI and HindII. Single stranded phage DNA was isolated and hybridized to a
5 chemically synthesized nucleotide with the sequence
3' TTTGCGGTAGTTGCCAGCTGTACCTTTATTGAGCATGA 5'.

The phage DNA was rendered partially double-stranded by simultaneous hybridization to plasmid DNA of M13mpl8 cleaved with BamHI and HindII.
10 After hybridization gaps were filled in by incubation with the Klenow fragment of DNA polymerase I (Boehringer) in a buffer containing all four deoxyribonucleotides. E. coli JM101 was transformed with the resulting plasmid DNA and correct colonies were picked up by hybridization to the
15 oligonucleotide which was end-labelled by gamma-(32)P-ATP and T4 polynucleotide kinase. Plasmid DNA was isolated and the mutagenized 0.6kb SalI-HindII fragment was isolated. It was ligated to the 2.3kb HindII-EcoRV fragment containing most of the 80 kDa coding sequence from pGB852 and the 9.2kb SalI-
20 EcoRV fragment containing the pUC sequence from pGB850. The resulting 80 kDa expression plasmid pGB853 was isolated (figure 16).

Example 14

25

CONSTRUCTION OF K. LACTIS VECTORS EXPRESSING A FUSION BETWEEN THE 92 kDa AND 80 kDa PROTEINS

In the fusion vector the PstI site at position 2660 of the
30 factor VIII sequence was ligated to the BamHI site at position 4749 as follows:

The 2.6kb SalI-PstI fragment encoding the 92 kDa protein from pCLB88, the 3.6kb BamHI-HindII fragment from pGB852, pUC19 digested with SalI and HindIII and an oligonucleotide with the
35 sequence ACGTCTAG were ligated giving rise to plasmid pGB860. A derivative of pGB850 was made in which the SalI site in the

polylinker was removed by a partial digestion with Sali followed by a 1 minute incubation with Bal31 as recommended by the manufacturer (BRL) yielding pGB850 dHSX. Next the factor VIII DNA encoding the fusion protein was put behind the
 5 lactase promoter by ligating the 5.8kb Sali-EcoRV fragment from pGB860 to the 92kb Sali-EcoRV fragment from pGB850 dHSX, resulting in the expression vector pGB861 (figure 17).

A vector for secretion of factor VIII subunits was obtained by replacing the leader for secretion of factor VIII
 10 in pGB 861 by a synthetic leader sequence as follows: the 1.9kb Sali-BamHI fragment encoding part of the 92 kDa protein of pGB861 was recloned in pTZ18R (United States Biochemical Corporation). A MluI site was introduced in the DNA at position 2 and 3 of the factor VIII protein sequence by using
 15 an oligonucleotide of the sequence
 5' ACCCAGGTAGTATCTacgcgtGGCACTAAAGCAGAA 3'.

The oligonucleotide was hybridized to the single stranded phage DNA of the plasmid containing the Sali-BamHI fragment in pTZ18R. After hybridization the gaps were filled
 20 in, and plasmids containing the oligonucleotide isolated as described in Example 10. From one of the correct plasmids the mutagenized 1.7kb Sali-BglII fragment was used to replace the corresponding fragment from pGB861, giving rise to pGB862. Plasmid pGB863 was cleaved with Sali and MluI. A synthetic DNA
 25 fragment was made encoding the synthetic leader sequence:

5' TCGACATGGCTTTCAGATCCTTGTTGGCTTTGTCCGGTTT...
 3' GTACCGAAAGTCTAGGAACAACCGAAACAGGCCAAA...

30 ...GTCCTGTGGTGCTTTGGCTGCTA3'
 ...CAGGACACCACGAAACCGACGATCGCG5'

It was ligated into pGB863 cleaved with Sali and BamHI giving rise to plasmid pGB864.

Example 15

TRANSFORMATION OF KLUYVEROMYCES LACTIS AND ANALYSIS OF THE RNA AND PROTEIN PRODUCTS OF THE TRANSFORMANTS.

5

All expression plasmids were linearized at the unique SacII restriction site and K. lactis was transformed according to Ito et al. (1984) using 0.2M lithium acetate, and 25 micrograms/ml of G418. Transformants were analysed by
10 isolation of chromosomal DNA followed by digestion with various restriction enzymes. Cell lines containing intact copies of the plasmid DNA integrated in the chromosome were grown up for 3 days in YEPD medium at 30°C for RNA and protein analysis.

15

For RNA analysis cells were harvested for 10 min at 5000 rpm. The pellet was suspended to an OD_{610nm} of 300 in a buffer containing 100 mM Tris-HCl, pH7.5; 100 mM LiCl; 0.1 mM EDTA and 0.5 mg/ml of heparin. To 1 ml of cells 2 g of glass beads (diameter 0.25-0.30 mm) were added and the
20 suspension was vortexed 5 times 30 seconds. Glass beads and cell debris were removed by centrifugation at 10,000 rpm for 5 min. The pellet was re-extracted with 0.5 ml of the above buffer and centrifuged. The supernatants were pooled, phenol extracted and ethanol precipitated. The RNA was incubated with
25 glyoxal, (McMaster and Carmichael, 1977), run on an agarose gel and blotted to Gene Screen Plus (NEN Research Products) as described by NEN. The result is shown in Figure 16. It is clear that both in K. lactis transformed with pGB850 or pGB852, RNA of the expected size can be found.

30

For protein analysis cells were harvested by a low speed centrifugation and suspended in 0.9% NaCl; 1mM phenylmethylsulfonyl fluoride (PMSF) (and 0.05% Tween 80 or 2% Nonidet P40 when indicated) to a final optical density of 500 at 610 nm. In a 10 ml conical test tube 1.2 g of glass beads
35 (diameter 0.25-0.30 mm) were added to 0.6 ml of the yeast cell suspension and the suspension was vortexed for 1 minute. The

cell debris and glass beads were removed by a low speed centrifugation. Samples were prepared and run on a SDS-polyacrylamide gel, blotted onto a nitrocellulose filter and incubated with antibodies as described in the General Methods 5 section.

Figure 19A shows the detection of the lactase-80 kDa fusion protein produced by K. lactis cells transformed with pGB852 as shown with the polyclonal antiserum RH 63275 and donkey anti-rabbit peroxidase (Promega Biotec). A band 10 corresponding to a protein product of the expected size (about 100 kDa) can be seen in lane 2 and, although much weaker, also in lane 4. In the parental K. lactis strain (lanes 1 and 3) this band cannot be detected.

The same results were obtained in a parallel 15 experiment where a blot with the same samples was incubated with polyclonal antiserum RH 63272 and goat anti-rabbit alkaline phosphatase from Promega Biotec (Figure 19B).

In a third experiment the presence of the fusion product could be shown by using monoclonal antiserum CLB- 20 CAG 65 and goat anti-mouse alkaline phosphatase (Figure 19C). In lanes 6-9 four independent K. lactis clones transformed with plasmid pGB852 are analysed each producing the lactase-80 kDa fusion product. They clearly possess a protein product of the expected size (approximately 100 kDa) not seen in the 25 parental yeast strain (lane 9) nor in K. lactis transformed with plasmid pGB850 containing the DNA sequence for the 92 kDa subunit of factor VIII (lanes 1-4).

For the detection of the 92 kDa-prochymosin fusion commercially available polyclonal antibodies (Chr. Hansen's 30 laboratory) against chymosin were used (Figure 19D).

Lanes 2-4 show the pattern of three independently obtained transformants of K. lactis with pGB851. A protein product of approximately the expected size (110 kDa) can be seen. After treatment of the cell extracts at pH 2 (General Methods, 4) 35 still most of the immuno-reacting material is located at the same position (lanes 10-12) but also two minor bands at the

approximate position of chymosin are visualised. So the fusion product is only partially processed by the pH 2 treatment. In the milk clotting assay chymosin activity (0.1 mg/l) is found in contrast to the parental yeast strain and the other 5 controls, demonstrating the presence of activated chymosin.

In another series of experiments it was shown that K. lactis also expresses 80 kDa and 92 kDa subunit proteins separately. In Fig. 19E a cell lysate of K. lactis transformed with pGB 853, run on a SDS polyacrylamide gel and blotted to a 10 nitrocellulose filter is shown. The blot was incubated with RH 63272 and goat anti-rabbit alkaline phosphatase. In Fig. 19F a similar experiment in which K. lactis was transformed with pGB 850 and monoclonal CLB-CAg 9 was used after blotting, is shown. Moreover, the two subunit proteins can be expressed 15 simultaneously in one cell (Fig. 19G). For this experiment pGB 850 and pGB 853 were cut with Sac II, and, after phenolization, mixed in equimolar amounts, and ligated. K. lactis was transformed with a mixture in which more than 50% of the DNA was in dimers. After growing up, cells were 20 lysed and strains expressing both subunits selected by immunoblotting and incubation with RH 63272 and CLB-CAg-9.

Fig. 19H shows that K. lactis is also able to express fusions between 92 kDa and 80 kDa proteins. Cells were transformed with pGB 861, lysed, proteins run on a SDS 25 polyacrylamide gel and analyzed with RH 63272. In most transformants a fusion protein of the expected size was found.

Secretion of a fusion protein between 92 kDa and 80 kDa subunits is demonstrated in Fig. 19I. K. lactis was 30 transformed with pGB864 and intra- and extracellular proteins determined separately.

It can be concluded that K. lactis can synthesize as intact products both the 80 kDa and 92 kDa proteins separately as well as fused to one another both intra- and 35 extracellularly.

In accordance with the subject invention, a number of factor VIII related proteins may be prepared by transformed microorganisms, including the 80 kDa and 92 kDa subunit proteins, a fusion protein of the 80 kDa and 92 kDa subunit proteins joined by a truncated bridge and fusion proteins of the subunit with an endogenous protein, where the endogenous protein is the N-terminus. The above products may be cytoplasmic or secreted in the medium. Individual products may be produced in the same or different cells and in the same or different cellular hosts.

It is evident from the above results, that the subject invention provides for products and methods which produce biologically and physiologically active proteins having factor VIII activity or being able to compete with factor VIII for antibodies which inhibit such activity. Thus, methods are provided using microbial hosts, with their inherent economies and efficiencies for the production of a complex protein, so as to provide drugs which can be used in place of the natural drug, so as to provide a stable, uniform, safe supply of factor VIII related products.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

The factor VIII cDNA insert of pCLB89. The amino acid sequence Ala-1 through Tyr-2332 of factor VIII and its signal sequence Met(-19) through Ser(-1) with the corresponding sequence of nucleotides. Phe-973 is encoded by the codon TTC.

0253455

10 20 30 40 50 60
GTCGACATGCAAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCATTCTGCTTT
-19 M Q I E L S T C F F L C L L R F C F -2

70 80 90 100 110 120
AGTGCCACCAGAAAGATACTACCTGGGTGCAGTGGAACTGTCATGGGACTATATGCAAAAGT
S A T R R Y Y L G A V E L S W D Y M Q S 19

130 140 150 160 170 180
GATCTCGGTGAGCTGCCTGTGGACGCAAGATTTCTCCTAGAGTGCCAAAATCTTTTCCA
D L G E L P V D A R F P P R V P K S F P

190 200 210 220 230 240
TTCAACACCTCAGTCGTGTACAAAAAGACTCTGTTTGTAGAATTCACGGATCACCTTTTC
F N T S V V Y K K T L F V E F T D H L F 59

250 260 270 280 290 300
AACATCGCTAAGCCAAGGCCACCCCTGGATGGGTCTGCTAGGTCCTACCATCCAGGCTGAG
N I A K P R P P W M G L L G P T I Q A E

310 320 330 340 350 360
GTTTATGATACAGTGGTCATTACACTTAAGAACATGGCTTCCCATCCTGTCACTCTTCAT
V Y D T V V I T L K N M A S H P V S L H 99

370 380 390 400 410 420
GCTGTTGGTGTATCCTACTGGAAGGCTTCTGAGGGAGCTGAATATGATGATCAGACCACT
A V G V S Y W K A S E G A E Y D D Q T S

430 440 450 460 470 480
CAAAGGGAGAAAGAAAGATGATAAAGTCTTCCCTGGTGGAAAGCCATACATATGTCTGGCAG
Q R E K E D D K V F P G G S H T Y V W Q 139

490 500 510 520 530 540
GTCCTGAAAGAGAAATGGTCCAATGGCCTCTGACCCACTGTGCCTTACCTACTCATATCTT
V L K E N G P M A S D P L C L T Y S Y L

550 560 570 580 590 600
TCTCATGTGGACCTGGTAAAGACTTGAATTCAGGCCTCATTGGAGCCCTACTAGTATGT
S H V D L V K D L N S G L I G A L L V C 179

610 620 630 640 650 660
AGAGAAGGGAGTCTGGCCAAGGAAAAGACACAGACCTTGACACAAATTTATACTACTTTTT
R E G S L A K E K T Q T L H K F I L L F

670 680 690 700 710 720
GCTGTATTTGATGAAGGGAAAAGTTGGCACTCAGAAACAAAGAACTCCTTGATGCAGGAT
A V F D E G K S W H S E T K N S L M Q D 219

730 740 750 760 770 780
AGGGATGCTGCATCTGCTCGGGCCTGGCCTAAAATGCACACAGTCAATGGTTATGTAAAC
R D A A S A R A W P K M H T V N B Y V N

790 800 810 820 830 840
AGGTCTCTGCCAGGTCTGATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTGGA
R S L P G L I G C H R K S V Y W H V I G 259

850 860 870 880 890 900
ATGGGCACCACTCCTGAAGTGCCTCAATATTCCTCGAAGGTCACACATTTCTTGTGAGG
M G T T P E V H S I F L E G H T F L V R

0253455

910 920 930 940 950 960
 AACCATCGCCAGGCGTCCTTGGAAATCTCGCCAATAACTTTCTTACTGCTCAAACTCT
 N H R Q A S L E I S P I T F L T A Q T L 299

970 980 990 1000 1010 1020
 TTGATGGACCTTGGACAGTTTCTACTGTTTTGTCATATCTCTTCCCACCAACATGATGGC
 L M D L G Q F L L F C H I S S H Q H D G

1030 1040 1050 1060 1070 1080
 ATGGAAGCTTATGTCAAAGTAGACAGCTGTCCAGAGGAACCCCACTACGAATGAAAAAT
 M E A Y V K V D S C P E E P Q L R M K N 339

1090 1100 1110 1120 1130 1140
 AATGAAGAAGCGGAAGACTATGATGATGATCTTACTGATTCTGAAATGGATGTGGTCAGG
 N E E A E D Y D D D L T D S E M D V V R

1150 1160 1170 1180 1190 1200
 TTTGATGATGACAACTCTCCTTCCTTTATCCAAATTCGCTCAGTTGCCAAGAAGCATCCT
 F D D D N S P S F I Q I R S V A K K H P 379

1210 1220 1230 1240 1250 1260
 AAACTTGGGTACATTACATTGCTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTC
 K T W V H Y I A A E E E D W D Y A P L V

1270 1280 1290 1300 1310 1320
 CTCGCCCCCGATGACAGAAAGTTATAAAAGTCAATATTTGAACAATGGCCCTCAGCGGATT
 L A P D D R S Y K S Q Y L N N G P Q R I 419

1330 1340 1350 1360 1370 1380
 GGTAGGAAGTACAAAAAGTCCGATTTATGGCATAACAGATGAAACCTTTAAGACTCGT
 G R K Y K K V R F M A Y T D E T F K T R

1390 1400 1410 1420 1430 1440
 GAAGCTATTGAGCATGAATCAGGAATCTTGGGACCTTTACTTTATGGGGAAAGTTGGAGAC
 E A I Q H E S G I L G P L L Y G E V G D 459

1450 1460 1470 1480 1490 1500
 AACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATATAACATCTACCCTCACGGA
 T L L I I F K N Q A S R P Y N I Y P H G

1510 1520 1530 1540 1550 1560
 ATCACTGATGTCCGTCCTTTGTATTCAAGGAGATTACCAAAGGTGTAAAACATTTGAAG
 I T D V R P L Y S R R L P K G V K H L K 499

1570 1580 1590 1600 1610 1620
 GATTTTCCAATTCTGCCAGGAGAAATATTCAAATATAAATGGACAGTGACTGTAGAAGAT
 D F P I L P G E I F K Y K W T V T V E D

1630 1640 1650 1660 1670 1680
 GGGCCAATAAATCAGATCCTCGGTGCCTGACCCGCTATTACTCTAGTTTTCGTTAATATG
 G P T K S D P R C L T R Y Y S S F V N M 539

1690 1700 1710 1720 1730 1740
 GAGAGAGATCTAGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATCTGTA
 E R D L A S G L I G P L L I C Y K E S V

1750 1760 1770 1780 1790 1800
 GATCAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGTATTT
 D Q R G N Q I M S D K R N V I L F S V F 579

0253455

1810 1820 1830 1840 1850 1860
GATGAGAACCGAAGCTGGTACCTCACAGAGAATATACACGCTTTCTCCCCAATCCAGCT
D E N R S W Y L T E N I Q R F L P N P A

1870 1880 1890 1900 1910 1920
GGAGTGACGCTTGAGGATCCAGAGTTCCAAGCCTCCAACATCATGCACAGCATCAATGGC
G V Q L E D P E F Q A S N I M H S I N G 619

1930 1940 1950 1960 1970 1980
TATGTTTTTGATAGTTTGCABTTGTGAGTTTGTTCATGAGGTGGCATACTGGTACATT
Y V F D S L Q L S V C L H E V A Y W Y I

1990 2000 2010 2020 2030 2040
CTAAGCATTGGAGCACAGACTGACTTCCTTTCTGTCTTCTCTCTGGATATACCTTCAA
L S I G A Q T D F L S V F F S G Y T F K 659

2050 2060 2070 2080 2090 2100
CACAAAATGGTCTATGAAGACACACTCACCTATTCCCATTCTCAGGAGAACTGTCTTC
H K M V Y E D T L T L F P F S G E T V F

2110 2120 2130 2140 2150 2160
ATGTCGATGGAAAACCCAGGTCTATGGATTCTGGGGTGCCACAACCTCAGACTTTTCGGAAC
M S M E N P G L W I L G C H N S D F R N 699

2170 2180 2190 2200 2210 2220
AGAGGCATGACCGCCTTACTGAAGGTTTCTAGTTGTGACAAGAACACTGGTGTATTATTAC
R G M T A L L K V S S C D K N T G D Y Y
12

2230 2240 2250 2260 2270 2280
GAGGACAGTTATGAAGATATTTCAACATACTTGCTGAGTAAAAACAATGCCATTGAACCA
E D S Y E D I S A Y L L S K N N A I E P 739

2290 2300 2310 2320 2330 2340
AGAAGCTTCTCCCAGAATTCAAGACACCCTAGCACTAGGCCAAAAGCAATTTAATGCCACC
R S F S Q N S R H P S T R Q K Q F N A T

2350 2360 2370 2380 2390 2400
ACAATTCCAGAAAATGACATAGAGAAGACTGACCCCTTGGTTTGCACACAGAACACCTATG
T I P E N D I E K T D P W F A H R T P M 779

2410 2420 2430 2440 2450 2460
CCTAAAATACAAAATGTCTCCTCTAGTGATTTGTTGATGCTCTTGCGACAGAGTCTACT
P K I Q N V S S S D L L M L L R Q S P T

2470 2480 2490 2500 2510 2520
CCACATGGGCTATCCTTATCTGATCTCCAAGAAGCCAAATATGAGACTTTTTCTGATGAT
P H G L S L S D L Q E A K Y E T F S D D 819

2530 2540 2550 2560 2570 2580
CCATCACCTGGAGCAATAGACAGTAATAACAGCCTGTCTGAAATGACACACTTCAGGCCA
P S P G A I D S N N S L S E M T H F R P

2590 2600 2610 2620 2630 2640
CAGCTCCATCACAGTGGGGACATGGTATTTACCCCTGAGTCAGGCCTCCAATTAAGATTA
Q L H H S G D M V F T P E S G L Q L R L 859

2650 2660 2670 2680 2690 2700
AATGAGAAACTGGGGACAACCTGCAGCAACAGAGTTGAAGAAACTTGATTTCAGAGTTTCT
N E K L G T T A A T E L K K L D F K V S

2710 2720 2730 2740 2750 2760
 AGTACATCAAATAATCTGATTTCAACAATTCATCAGACAATTTGGCAGCAGGTACTGAT
 S T S N N L I S T I P S D N L A A G T D 899

2770 2780 2790 2800 2810 2820
 AATACAAGTTCCTTAGGACCCCCAAGTATGCCAGTTCATTATGATAGTCAATTAGATACC
 N T S S L G P P S M P V H Y D S Q L D T

2830 2840 2850 2860 2870 2880
 ACTCTATTTGGCAAAAAGTCATCTCCCCTTACTGAGTCTGGTGGACCTCTGAGCTTGAGT
 T L F G K K S S P L T E S G G P L S L S 939

2890 2900 2910 2920 2930 2940
 GAAGAAAATAATGATTCAAAGTTGTTAGAATCAGGTTTAATGAATAGCCAAGAAAGTTCA
 E E N N D S K L L E S G L M N S Q E S S

2950 2960 2970 2980 2990 3000
 TGGGGAAAAAATGTATCGTCAACAGAGAGTGGTAGGTTATTCAAAGGGAAAAGAGCTCAT
 W G K N V S S T E S G R L F K G K R A H 979

3010 3020 3030 3040 3050 3060
 GGACCTGCTTTGTTGACTAAAGATAATGCCTTATTCAAAGTTAGCATCTCTTTGTTAAAG
 G P A L L T K D N A L F K V S I S L L K

3070 3080 3090 3100 3110 3120
 ACAAACAAACTTCCAATAATTCAGCAACTAATAGAAAGACTCACATTGATGGCCCATCA
 T N K T S N N S A T N R K T H I D G P S 1019

3130 3140 3150 3160 3170 3180
 TTATTAATTGAGAATAGTCCATCAGTCTGGCAAAATATATTAGAAAGTGACACTGAGTTT
 L L I E N S P S V W Q N I L E S D T E F

3190 3200 3210 3220 3230 3240
 AAAAAAGTGACACCTTTGATTCATGACAGAATGCTTATGGACAAAAATGCTACAGCTTTG
 K K V T P L I H D R M L M D K N A T A L 1059

3250 3260 3270 3280 3290 3300
 AGGCTAAATCATATGTCAAATAAACTACTTCATCAAAAAACATGGAAATGGTCCAACAG
 R L N H M S N K T T S S K N M E M V Q Q

3310 3320 3330 3340 3350 3360
 AAAAAAGAGGGCCCCATTCCACCAGATGCACAAAATCCAGATATGTCGTTCTTTAAGATG
 K K E G P I P P D A Q N P D M S F F K M 1099

3370 3380 3390 3400 3410 3420
 CTATTCTTGCCAGAATCAGCAAGGTGGATACAAAGGACTCATGGAAAGAACTCTCTGAAC
 L F L P E S A R W I Q R T H G K N S L N

3430 3440 3450 3460 3470 3480
 TCTGGGCAAGGCCCCAGTCCAAAGCAATTAGTATCCTTAGGACCAGAAAAATCTGTGGAA
 S G Q G P S P K Q L V S L G P E K S V E 1139

3490 3500 3510 3520 3530 3540
 GGTGAGAATTTCTTGTCTGAGAAAAACAAAGTGGTAGTAGGAAAGGGTGAATTTACAAAG
 G Q N F L S E K N K V V V G K G E F T K

3550 3560 3570 3580 3590 3600
 GACGTAGGACTCAAAGAGATGGTTTTTCCAAGCAGCAGAAACCTATTTCTTACTAACTTG
 D V G L K E M V F P S S R N L F L T N L 1179

3610 3620 3630 3640 3650 3660
GATAATTTACATGAAAATAATACACACAATCAAGAAAAAAATTCAGGAAGAAATAGAA
D N L H E N N T H N Q E K K I Q E E I E

3670 3680 3690 3700 3710 3720
AAGAAGGAAACATTAATCCAAGAGAATGTAGTTTTGCCTCAGATACATACAGTGACTGGC
K K E T L I Q E N V V L P Q I H T V T G 1219

3730 3740 3750 3760 3770 3780
ACTAAGAATTTTCATGAAGAACCTTTTCTTACTGAGCACTAGGCAAAATGTAGAAGGTTCA
T K N F M K N L F L L S T R Q N V E G B

3790 3800 3810 3820 3830 3840
TATGACGGGGCATATGCTCCAGTACTTCAAGATTTTAGGTCATTAAATBATTCAACAAAT
Y D G A Y A P V L Q D F R S L N D S T N 1259

3850 3860 3870 3880 3890 3900
AGAACAAAGAAACACACAGCTCATTCTCAAAAAAAGGGGAGGAAGAAACCTTGGAAAGGC
R T K K H T A H F S K K G E E E N L E G

3910 3920 3930 3940 3950 3960
TTGGGAAATCAAACCAAGCAAATTGTAGAGAAATATGCATGCACCACAAGGATATCTCCT
L G N Q T K Q I V E K Y A C T T R I S P 1299

3970 3980 3990 4000 4010 4020
AATACAAGCCAGCAGAATTTTGTACGCAACGTAGTAAGAGAGCTTTGAAACAATTCAGA
N T S Q Q N F V T Q R S K R A L K Q F R

4030 4040 4050 4060 4070 4080
CTCCCACTAGAAGAAACAGAACTTGAAAAAAGGATAATTGTGGATGACACCTCAACCCAG
L P L E E T E L E K R I I V D D T S T Q 1339

4090 4100 4110 4120 4130 4140
TGGTCCAAAAACATGAAACATTTGACCCCGAGCACCCTCACACAGATAGACTACAATGAG
W S K N M K H L T P S T L T Q I D Y N E

4150 4160 4170 4180 4190 4200
AAGGAGAAAGGGGCCATTACTCAGTCTCCCTTATCAGATTGCCTTACGAGGAGTCATAGC
K E K G A I T Q S P L S D C L T R S H S 1379

4210 4220 4230 4240 4250 4260
ATCCCTCAAGCAAATAGATCTCCATTACCCATTGCAAAGGTATCATCATTTCCATCTATT
I P Q A N R S P L P I A K V S S F P S I

4270 4280 4290 4300 4310 4320
AGACCTATATATCTGACCAGGGTCCTATTCCAAGACAACCTCTTCTCATCTTCCAGCA
R P I Y L T R V L F Q D N S S H L P A A 1419

4330 4340 4350 4360 4370 4380
TCTTATAGAAAGAAAGATTCTGGGGTCCAAGAAAGCAGTCATTTCTTACAAGGAGCCAAA
S Y R K K D S G V Q E S S H F L Q G A K

4390 4400 4410 4420 4430 4440
AAAAATAACCTTTCTTTAGCCATTCTAACCTTGGAGATGACTGGTGATCAAGAGAGGTT
K N N L S L A I L T L E M T G D Q R E V 1459

4450 4460 4470 4480 4490 4500
GGCTCCCTGGGGACAAGTGCCACAAATTCAGTCACATACAAGAAAGTTGABAACACTGTT
G S L G T S A T N S V T Y K K V E N T V

4510 4520 4530 4540 4550 4560
 CTCCCGAAACCAGACTTGCCCAAAACATCTGGCAAAGTTGAATTGCTTCCAAAAGTTCAC
 L P K P D L P K T S G K V E L L P K V H 1499

4570 4580 4590 4600 4610 4620
 ATTTATCAGAAGGACCTATTCCCTACGGAACTAGCAATGGGTCTCCTGGCCATCTGGAT
 I Y Q K D L F P T E T S N G S P G H L D

4630 4640 4650 4660 4670 4680
 CTCGTGGAAAGGGAGCCTTCTTCAGGGAAACAGAGGGAGCGATTAAAGTGGGAATGAAGCAAAC
 L V E G S L L G G T E G A I K W N E A N 1539

4690 4700 4710 4720 4730 4740
 AGACCTGGAAAAGTTCCCTTTCTGAGAGTAGCAACAGAAAGCTCTGCAAAGACTCCCTCC
 R P G K V P F L R V A T E S S A K T P S

4750 4760 4770 4780 4790 4800
 AAGCTATTGGATCCTCTTGCTTGGGATAACCACTATGGTACTCAGATACCAAAAAGAAGAG
 K L L D P L A W D N H Y G T Q I P K E E 1579

4810 4820 4830 4840 4850 4860
 TGGAAATCCCAAGAGAAGTCACCAGAAAAACAGCTTTTAAGAAAAAGGATACCATTTTG
 W K S Q E K S P E K T A F K K K D T I L

4870 4880 4890 4900 4910 4920
 TCCCTGAACGCTTGTGAAAGCAATCATGCAATAGCAGCAATAAATGAGGGACAAAATAAG
 S L N A C E S N H A I A A I N E G Q N K 1619

4930 4940 4950 4960 4970 4980
 CCCGAAATAGAAGTCACCTGGGCAAAGCAAGGTAGGACTGAAAGGCTGTGCTCTCAAAC
 P E I E V T W A K Q G R T E R L C S Q N
 37 38

4990 5000 5010 5020 5030 5040
 CCACCAGTCTTGAAACGCCATCAACGGGAAATAACTCGTACTACTCTTCAGTCAGATCAA
 P P V L K R H Q R E I T R T T L Q S D Q 1659
 49

5050 5060 5070 5080 5090 5100
 GAGGAAATTGACTATGATGATACCATATCAGTTGAAATGAAGAAAGGAGATTTTGACATT
 E E I D Y D D T I S V E M K K E D F D I

5110 5120 5130 5140 5150 5160
 TATGATGAGGATGAAAATCAGAGCCCCCGCAGCTTTCAAAGAAAACACGACACTATTTT
 Y D E D E N Q S P R S F Q K K T R H Y F 1699
 89 90

5170 5180 5190 5200 5210 5220
 ATTGCTGCAGTGGAGAGGCTCTGGGATTATGGGATGAGTAGCTCCCCACATGTTCTAAGA
 I A A V E R L W D Y G M S S S P H V L R

5230 5240 5250 5260 5270 5280
 AACAGGGCTCAGAGTGGCAGTGTCCCTCAGTTCAAGAAAGTTGTTTTCCAGGAATTTACT
 N R A Q S G S V P Q F K K V V F Q E F T 1739

5290 5300 5310 5320 5330 5340
 GATGGCTCCTTTACTCAGCCCTTATACCGTGGAGAACTAAATGAACATTTGGGACTCCTG
 D G S F T Q P L Y R G E L N E H L G L L

5350 5360 5370 5380 5390 5400
 GGGCCATATATAAGAGCAGAAGTTGAAGATAATATCATGGTAACCTTCAGAAATCAGGCC
 G P Y I R A E V E D N I M V T F R N Q A 1779

5410 5420 5430 5440 5450 5460
TCTCGTCCCTATTCTTCTATTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGA
S R P Y S F Y S S L I S Y E E D Q R Q G

5470 5480 5490 5500 5510 5520
GCAGAACCTAGAAAAAAGCTTTGTCAAGCCTAATGAAACCAAACTTACTTTTGGAAAGTG
A E P R K N F V K P N E T K T Y F W K V 1819

5530 5540 5550 5560 5570 5580
CAACATCATATGGCACCCTAAAGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCTCT
Q H H M A P T K D E F D C K A W A Y F S

5590 5600 5610 5620 5630 5640
GATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCTGGTCTGCCAC
D V D L E K D V H S G L I G P L L V C H 1859

5650 5660 5670 5680 5690 5700
ACTAACACACTGAACCCTGCTCATGGGAGACAAGTGACAGTACAGGAATTTGCTCTGTTT
T N T L N P A H G R Q V T V Q E F A L F

5710 5720 5730 5740 5750 5760
TTCACCATCTTTGATGAGACCAAAAGCTGGTACTTCACTGAAAATATGGAAAGAACTGC
F T I F D E T K S W Y F T E N M E R N C 1899

5770 5780 5790 5800 5810 5820
AGGGCTCCCTGCAATATCCAGATGGAAGATCCCCTTTTAAAGAGAATTATCGCTTCCAT
R A P C N I Q M E D P T F K E N Y R F H

5830 5840 5850 5860 5870 5880
GCAATCAATGGCTACATAATGGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGG
A I N G Y I M D T L P G L V M A Q D Q R 1939

5890 5900 5910 5920 5930 5940
ATTCGATGGTATCTGCTCAGCATGGGCAGCAATGAAAACATCCATTCTATTTCATTTCAGT
I R W Y L L S M G S N E N I H S I H F S

5950 5960 5970 5980 5990 6000
GGACATGTGTTCACTGTACGAAAAAAGAGGAGTATAAAATGGCACTGTACAATCTCTAT
G H V F T V R K K E E Y K M A L Y N L Y 1979

6010 6020 6030 6040 6050 6060
CCAGGTGTTTTTGAGACAGTGGAAATGTTACCATCCAAAGCTGGAATTTGGCGGGTGGAA
P G V F E T V E M L P S K A G I W R V E

6070 6080 6090 6100 6110 6120
TGCCTTATTGGCGAGCATCTACATGCTGGGATGAGCACACTTTTCTGGTGTACAGCAAT
C L I G E H L H A G M S T L F L V Y S N 2019

6130 6140 6150 6160 6170 6180
AAGTGTCAAGTCCCTGGGAATGGCTTCTGGACACATTAGAGATTTTCAGATTACAGCT
K C Q T P L G M A S G H I R D F Q I T A

6190 6200 6210 6220 6230 6240
TCAGGACAATATGGACAGTGGGCCCCAAAGCTGGCCAGACTTCATTATTCCGGATCAATC
S G Q Y G Q W A P K L A R L H Y S G S I 2059

6250 6260 6270 6280 6290 6300
AATGCCTGGAGCACCAAGGAGCCCTTTTCTTGGATCAAGGTGGATCTGTTGGCACCAATG
N A W S T K E P F S W I K V D L L A P M

6310 6320 6330 6340 6350 6360
ATTATTACGGCATCAAGACCCAGGGTGCCCGTCAGAAGTTCTCCAGCCTCTACATCTCT
I I H G I K T Q G A R Q K F S S L Y I S 2099

6370 6380 6390 6400 6410 6420
CAGTTTATCATCATGTATAGTCTTGATGGGAAGAAAGTGGCAGACTTATCGAGGAAATTCC
Q F I I M Y S L D G K K W Q T Y R G N S

6430 6440 6450 6460 6470 6480
ACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTTCATCTGGGATAAAACACAATATT
T G T L M V F F G N V D S S G I K H N I 2139

6490 6500 6510 6520 6530 6540
TTTAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAACTCATTATAGCATTGCG
F N P P I I A R Y I R L H P T H Y S I R

6550 6560 6570 6580 6590 6600
AGCACTCTTCGCATGGAGTGGATGGGCTGTGATTTAAATAGTTGCAGCATGCCATTGGGA
S T L R M E W M G C D L N S C S M P L G 2179

6610 6620 6630 6640 6650 6660
ATGGAGAGTAAAGCAATATCAGATGCACAGATTACTGCTTCATCCTACTTTACCAATATG
M E S K A I S D A Q I T A S S Y F T N M

6670 6680 6690 6700 6710 6720
TTTGCCACCTGGTCTCCTTCAAAAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTGG
F A T W S P S K A R L H L Q G R S N A W 2219

6730 6740 6750 6760 6770 6780
AGACCTCAGGTGAATAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAA
R P Q V N N P K E W L Q V D F Q K T M K

6790 6800 6810 6820 6830 6840
GTCACAGGAGTAACTACTCAGGGAGTAAATCTCTGCTTACCAGCATGTATGTGAAGGAG
V T G V T T Q G V K S L L T S M Y V K E 2259

6850 6860 6870 6880 6890 6900
TTCCTCATCTCCAGCAGTCAAGATGGCCATCAGTGGACTCTCTTTTTTTCAGAAATGGCAAA
F L I S S S Q D G H Q W T L F F Q N G K

6910 6920 6930 6940 6950 6960
GTAAAGGTTTTTCAGGGAAATCAAGACTCCTTCACACCTGTGGTGAAGTCTCTAGACCCA
V K V F Q G N Q D S F T P V V N S L D P 2299

6970 6980 6990 7000 7010 7020
CCGTTACTGACTCGCTACCTTCGAATTCACCCCAAGAGTTGGGTGCACCAAGATTGCCCTG
P L L T R Y L R I H P Q S W V H Q I A L

7030 7040 7050 7060 7070 7080
AGGATGGAGGTTCTGGGCTGCGAGGCACAGGACCTCTACTGAGGGTGGCCACTGCAGCAC
R M E V L G C E A Q D L Y
2332 stop

7090 7100 7110 7120 7130 7140
CTGCCACTGCCGTCACCTCTCCCTCCTCAGCTCCAGGGCAGTGTCCCTCCCTGGCTTGCC

7150 7160 7170 7180 7190 7200
TTCTACCTTTGTGCTAAATCCTAGCAGACACTGCCTTGAGGCCTCCTGAATTAAGTATCA

7210 7220 7230 7240 7250 7260
TCAGTCCTGCATTTCTTTGGTGGGGGGCCAGGAGGGTGCATCCAATTTAACTTAACTCTT

7270 7280 7290 7300 7310 7320
ACCTATTTTCTGCAGCTGCTCCCAGATTACTCCTTCCTTCCAATATAACTAGGCCAAAAAG

7330 7340 7350 7360 7370 7380
AAGTGAGGAGAAACCTGCATGAAAGCATTCTTCCTGAAAAGTTAGGCCTCTCAGAGTCA

7390 7400 7410 7420 7430 7440
CCACTTCCTCTGTTGTAGAAAACTATGTGATGAACTTTGAAAAAGATATTTATGATGT

7450 7460 7470 7480 7490 7500
TAACATTTTCAGGTTAAGCCTCATACGTTTAAATAAACTCTCAGTTGTTTATTATCCTG

7510 7520 7530 7540 7550 7560
ATCAAGCATGGAACAAAGCATGTTTCAGGATCAGATCAATACAATCTTGGAGTCAAAAGG

7570 7580 7590 7600 7610 7620
CAAATCATTTCGACAATCTGCAAAATGGAGAGAATACAATAACTACTACAGTAAAGTCTG

7630 7640 7650 7660 7670 7680
TTTCTGCTTCCTTACACATAGATATAATTATGTTATTTAGTCATTATGAGGGGCACATTC

7690 7700 7710 7720 7730 7740
TTATCTCCAAACTAGCATTCTTAACTGAGAATTATAGATGGGGTTCAAGAATCCCTAA

7750 7760 7770 7780 7790 7800
GTCCCCTGAAATTATATAAGGCATTCTGTATAAATGCAATGTGCATTTTCTGACGAGT

7810 7820 7830 7840 7850 7860
GTCCATAGATAAAAAGCCATTGTTGTTAATTCTGACCAATAAAAAATAAGTCAGGAGG

7870 7880 7890 7900 7910 7920
ATGCAATTGTTGAAAGCTTTGAAATAAAATAACAATGTCTTCTTGAAATTTGTGATGGCC

7930 7940 7950 7960 7970 7980
AAGAAAGAAAATGATGATGACATTAGGCTTCTAAAGGACATACATTTAATATTTCTGTGG

7990 8000 8010 8020 8030 8040
AAATATGAGGAAAATCCATGGTTATCTGAGATAGGAGATACAACTTTGTAATTCTAATA

8050 8060 8070 8080 8090 8100
ATGCACTCAGTTTACTCTCTCCCTCTACTAATTTCTGCTGAAAATAACACAACAAAAT

-53 k-

0253455

8110 8120 8130 8140 8150 8160
GTAACAGGGGAAATTATATACCGTGACTGAAAACAGAGTCCTACTTACATAGTTGAAAT

8170 8180 8190 8200 8210 8220
ATCAAGGAGGTCAGAAGAAAATTGGACTGGTGAAAACAGAAAAACACTCCAGTCTGCCA

8230 8240
TATCACCACACAATAGGATCC

REFERENCES

- Anagnostopoulos, C. and Spizizen, J. (1961) J. Bacteriol. 81, 741-746.
- 5 Andreoli, P.M. (1985) Mol. Gen. Genet. 199, 372-380.
- Ano, T. et al., (1986) Mol. Gen. Genet. 202, 416-420.
- 10 Arraj, J.A. and Marinus, M.G., (1983) J. Bacteriol, 153, 562-565
- Aviv, H. and Leder, P. (1972) Proc. Natl. Acad. Sci. USA 69, 1408-1412
- 15 Breunig, K.D. et al (1984) Nucl. Acids Res. 12, 2327-2341.
- Brinkhous, K.M. et al., (1985) Proc. Natl. Acad. Sci. USA 82, 8752-8756
- 20 Chang, A.C.Y. and Cohen, S.N. (1978) J. Bacteriol. 134, 1141-1156
- Chang, S. and Cohen, S.N. (1979) Mol. Gen. Genet. 168, 111-25 115.
- Chargwin, J.M. et al., (1979) Biochemistry 18, 5294-5299
- Das, S. and Hollenberg, C.P. (1982) Curr. Genet. 6, 123-30 128.
- Eaton, D. et al., (1986) Biochemistry 25, 505-512
- Enger-Valk, B.E. Ph.D. thesis (1981), University of 35 Wageningen, The Netherlands.

- Fay, P.J. et al., (1986) Biochim. Biophys. Acta 871, 268-278
- Foltman, B. (1970) Methods Enzymol. 19, 421-436
- 5 Galfré, G. et al., (1977) Nature 266, 550-552
- Gandhi, A.P. and Kjaergaard, L. (1975) Biotechnol. Bioeng. 17, 1109-1118.
- 10 Grunstein, M. and Hogness, D.S. (1975) Proc. Natl. Acad. Sci. USA 72 (1975) 3961-3965.
- Gubler, U. and Hoffman, B.J. (1983) Gene 25, 263-269.
- Hanahan, D. (1983) J. Mol. Biol. 166, 557-580.
- 15 Holmes, D.S. and Quigley, M. (1981) Anal. Biochem 114, 193-197.
- Ito, H., Murata, K. and Kimura, A. (1984) Agric. Biol. Chem.
- 20 48, 341-347.
- Kawamura, F. and Doi, R.H. (1984) J. Bacteriol. 160, 442-444
- Laemli, U.K. (1970) Nature 227, 680-685.
- 25 Maniatis, T. et al., (1982). Molecular cloning. Cold Spring Harbor Laboratory.
- Marinus, M.G. and Morris, N.R. (1973), J. Bacteriol. 114,
- 30 143-1150
- Matsumura, M. et al., (1984) J. Bacteriol. 160, 413-420.
- Maxam, A.M. and Gilbert, W. (1980) Methods Enzymol. 65, 499-
- 35 560.

- McMaster, G.K. and Carmichael, G.G. (1977) Proc. Natl. Acad. Sci. USA 74, 4835-4838.
- Melton, D.A. et al (1984) Nucl. Acids. Res. 12, 7035-7056.
- 5
Messing, J. and Vieira, J. (1982) Gene 19, 269-276.
- Meyer, H.P. and Fiechter, A. (1985) Appl. Environ. Microbiol. 50, 503-507.
- 10
Miller, J.H. (1972), Experiments in Molecular Genetics, Cold Spring, Harbor Laboratory, New York.
- Niaudet, B. et al., (1984) Mol. Gen. Genet. 197, 46-54.
- 15 Ohmura, K. et al., (1984) J. Biochem. 95, 87-93.
- Palva, I. et al., (1983) Gene 22, 229-235.
- Polak, J. and Novick, R.P. (1982) Plasmid 7 152-162.
- 20
Sanger, F. et al. (1977) Proc. Natl. Acad. Sci USA 74, 5463-5467.
- Sarvas, M. (1986), Current Topics in Microbiol. and Immunol. 25 103-125 H.C. Wu and P.C. Tai eds., Springer Verlag.
- Shiroza, T. et al. (1985) Gene 34, 1-8.
- Stel, H.V. (1984) Ph.D. Thesis, University of Amsterdam, The Netherlands.
- 30
Stel, H.V. et al., (1983) Nature 303, 530-533.
- Thorne, C.B. and Stull, H.B. (1966) J. Bacteriol. 91, 1012-1020.
- 35 Toole, J.J. et al (1984) Nature 312, 342-347.

Towbin H. et al., (1979) Proc. Natl. Acad. Sci. USA 76,
4350-4354.

Truett, M.A. et al., (1985) DNA 4, 333-349.

5 Twigg, A.J. and Sherratt, D. (1980) Nature 283, 216-218.

Ulmanen, I. et al., (1985) J. Bacteriol. 162, 176-182.

Van Mourik J. A. and Mochtar, J.A. (1970) Biochim.
10 Biophys. Acta 221, 677-679.

Vehar, G. et al. (1984) Nature 312, 337-342.

Wood, W.I. et al. (1984). Nature 312, 330-337.

15

Yanisch-Perron, C. et al., (1985) Gene 33, 103-119.

CLAIMS

1. An expression cassette comprising in the
5 direction of transcription, a transcriptional and
translational initiation regulatory region, optionally a
secretion signal sequence, an open reading frame encoding an
amino acid sequence immunologically cross-reactive with factor
VIII, and a translational and transcriptional termination
10 regulatory region, said regulatory regions being functional in
a microbial host.

2. An expression cassette according to claim 1,
wherein said open reading frame substantially encodes at least
15 one of the 80 kDa protein or the 92 kDa protein of factor
VIII.

3. An expression cassette according to claim 2,
wherein said open reading frame encodes the 80 kDa and 92 kDa
20 proteins joined by other means than the natural polypeptide
bridge of factor VIII.

4. A vector comprising a replication system
functional in a microbial host and an expression cassette
25 comprising in the direction of transcription, a
transcriptional and translational initiation regulatory
region, optionally a secretion signal sequence, an open
reading frame encoding an amino acid sequence immunologically
cross-reactive with factor VIII, and a translational and
30 transcriptional termination regulatory region, said regulatory
regions being functional in a microbial host.

5. A vector according to claim 4, wherein said
vector further comprises a marker for selection in a microbial
35 host.

6. A vector according to claim 4, wherein said

open reading frame substantially encodes at least one of the 80 kDa protein or the 92 kDa protein of factor VIII.

7. A vector according to claim 4, wherein said
5 open reading frame substantially encodes the 80 kDa protein.

8. A vector according to claim 4, wherein said
open reading frame substantially encodes the 92 kDa protein.

10 9. A vector according to claim 4, wherein said
open reading frame encodes the 80 kDa and 92 kDa proteins
joined by other means than the natural polypeptide bridge of
factor VIII.

15 10. A vector according to claim 9, wherein said 80
kDa and 92 kDa proteins are in the natural order of factor
VIII.

20 11. Plasmid pGB850

12. Plasmid pGB852

13. Plasmid pGB853

25 14. Plasmid pGB861

15. Plasmid pGB864

30 16. Plasmid pOL92

17. Plasmid pOL80

18. Plasmid pPROM92/80PB

35 19. Vector pOL5

20. The preparation of a microbial host, characterized in that a Bacillus subtilis or a Bacillus licheniformis, which is alpha-amylase deficient and/or exoprotease deficient is transformed with a vector according to any one of claims 4-10 or 16-18.

21. A transformed microbial host comprising an expression cassette according to any one of claims 1-3.

10 22. A microbial host according to claim 21, wherein said host is a bacterium or a yeast.

23. A microbial host according to claim 22, wherein said bacterium is a Bacillus and the yeast is a Kluyveromyces.

15 24. A microbial host according to claim 23, wherein said bacterium is Bacillus subtilis or Bacillus licheniformis.

25 25. A Bacillus host according to any one of claims 23-24 wherein said expression cassette is present on the plasmid pOL92 , pOL80 or pPROM92/80PB.

26. A microbial host according to claim 23, wherein said Kluyveromyces is Kluyveromyces lactis.

25 27. A Kluyveromyces host according to claim 23 or 26, wherein said expression cassette is present on the plasmid pGB850, pGB852, pGB853, pGB861 or pGB864.

30 28. A process for producing a polypeptide in isolatable form cross-reactive with factor VIII, said process comprising;

35 cultivating a microbial host according to any of claims 21-27 in a nutrient medium to produce said polypeptide; and isolating said polypeptide.

29. A process according to claim 28, wherein said open reading frame is joined at its 5' terminus to a DNA sequence encoding a signal sequence functional in said microbial host for secretion of said polypeptide and wherein
5 said signal sequence is processed.

30. A process according to claim 29, wherein said microbial host is a Kluyveromyces.

10 31. A process according to claim 29, wherein said microbial host is a Bacillus.

32. A pharmaceutical composition comprising a polypeptide produced by the process comprising:
15 cultivating a microbial host according to any of claims 21-27 in a nutrient medium to produce said polypeptide; and
isolating said polypeptide,
and a pharmaceutically acceptable carrier or diluent and
20 optionally an adjuvant.

33. A polypeptide produced by the process comprising: cultivating a microbial host according to any of claims 21-27 in a nutrient medium to produce said polypeptide, and
25 isolating said polypeptide.

34. An antibody to a protein obtained by a process according to any one of claims 28-31.

30 35. A monoclonal antibody according to claim 34.

36. A method of purifying protein having factor VIII activity which comprises bringing a liquid phase comprising protein and a content of impurities into contact with a solid
35 phase comprising antibody according to claim 34 or 35, separating the liquid phase from the solid phase and eluting

from the solid phase protein with a reduced content of impurities.

37. A method of diagnosing a blood sample for factor VIII abnormalities which comprises bringing the sample into contact with an antibody according to claim 34 or 35 and assaying for the presence or absence of a conjugate of the antibody with factor VIII antigen as a measure of the factor VIII protein in the sample.


1/27

0253455

Fig. 1



Fig. 2



3/27

0253455

m.c.s. EcoRI XmaIII XmaI Sall HindIII
5'pGAA TTC GCG GCC GCC CGG GTC GAC TCA AGC TTA AAT
TCG CGG CCG CCC GGG TCG ACT CAA GCT TAA ATT
CGC GGC CGC CCG GGT CGA CTC AAG CTT AAA 3'




Fig. 3

4/27

0253455

Fig. 4





Fig 5A



6/27

0253455





Fig. 5 B


Fig. 5C



0253455

8/27


Fig. 6



9/27

0253455


Fig. 7



0253455

10/27


Fig. 8



11/27

0253455

Fig. 9



12/27

0253455

1 2 3 4 5 6 7





Fig. 10 AB

A

1 2 3



4 5 6 7




B

13/27

0253455


1 2 3 4 5 6 7



C

Fig. 10 C D

1 2 3 4 5 6 7 8



D

14/27

0253455





Fig. 11

0253455

15/27

Fig. 12



16/27

0253455

Fig. 13






Fig. 14



0253455

10/27


Fig. 15



0253455


19/27

Fig. 16




20/27

Fig. 17



21/27


Fig. 18 A B



0253455

22/27


Fig. 19 AB



0253455

23/27


Fig. 19C



0253455

24/27

Fig. 19 D




0253455

25/27

Fig. 19 E

1 2 3 4 5 6




E

0253455

26/27


Fig 19 F G H



0253455

27/27

Fig 19 I





European Patent
Office

EUROPEAN SEARCH REPORT

0253455

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 87201379.2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
D,A	EP - A2 - 0 150 735 (CHIRON CORPORATION) * Pages 13,14; claims 1,4 * --	1-10, 20-24, 34,35	C 12 N 15/00 C 12 N 1/20 C 12 N 1/16 C 12 P 21/00 A 61 K 37/02 C 07 K 13/00 A 61 K 39/395 G 01 N 33/68 G 01 N 33/577
D,A	EP - A1 - 0 160 457 (GENENTECH, INC.) * Claims * --	1,4	(C 12 N 1/20 C 12 R 1:125) (C 12 N 1/20 C 12 R 1:10) (C 12 N 1/20 C 12 R 1:07)
D,A	DNA, vol. 4, no. 1, February 1985, New York M.A.TRUETT et al. "Characterization of the Polypeptide Composition of Human Factor VIII:C and the Nucleotide Sequence and Expression of the Human Kidney cDNA" pages 333-349 * Pages 333-335 * --	36,37	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 12 N C 12 P A 61 K C 07 K G 01 N
D,A	BIOCHEMICA ET BIOPHYSICA ACTA, vol. 871, June 23, 1986, Amsterdam P.J.FAY et al. "The size of human factor VIII heterodimers and the effects produced by thrombin" pages 268-278 * Totality * --		
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 30-09-1987	Examiner WOLF
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons a : member of the same patent family, corresponding document</p>			

0253455



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

-2-

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 87201379.2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
D,A	<p>BIOCHEMISTRY, vol. 25, no. 2, January 28, 1986, New York</p> <p>D.EATON et al. "Proteolytic Processing of Human Factor VIII. Correlation of Specific Cleavages by Thrombin, Factor Xa, and Activated Protein C with Activation and Inactivation of Factor VIII Coagulant Activity" pages 505-512</p> <p>* Totality *</p> <p>----</p>		
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 30-09-1987	Examiner WOLF
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			



**DECLARATION PURSUANT TO RULE 28, PARAGRAPH 4,
OF THE EUROPEAN PATENT CONVENTION**

The applicant has informed the European Patent Office that, until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, the availability of the micro-organism(s) identified below, referred to in paragraph 3 of Rule 28 of the European Patent Convention, shall be effected only by the issue of a sample to an expert.

IDENTIFICATION OF THE MICRO-ORGANISMS

Accession numbers of the deposits: CBS 304.86
CBS 303.86
CBS 305.86
CBS 306.86

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.